



CCDS

CROP CONDITION AND DEVELOPMENT STAGES

JANUARY 13, 2022



DEPARTAMENT & REGIONS

HEAD OF DEPARTMENT

Ing. Esteban J. Copati
ecopati@bc.org.ar

CROP ANALYST

Ing. Martin López
martinlopez@bc.org.ar

CROP ANALYST

Ing. Daniela A. Venturino
dventurino@bc.org.ar

CROP ANALYST


Ing. Andrés Paterniti
apaterniti@bc.org.ar

CROP SURVEY & ANALYSIS OF CROP CONDITION AND DEVELOPMENT STAGES

Justo Fernández Vidal
jfernandez@bc.org.ar

Jorgelina Mediate
jmediate@bc.org.ar

Joaquín Pellejero
jpellejero@bc.org.ar



I - NWA (North-West Argentina)
 II - NEA (North-East Argentina)
 III - North-Central Córdoba
 IV - South Córdoba
 V - North-Central Santa Fe
 VI - North Belt
 VII - South Belt
 VIII - East-Central Entre Ríos

IX - North La Pampa - West Buenos Aires
 X - Central Buenos Aires
 XI - South-West de Buenos Aires - South La Pampa
 XII - South-East Buenos Aires
 XIII - San Luis
 XIV - Cuenca del Salado
 XV - Others


CONTACT

Av. Corrientes 123
 C1043AAB - CABA
 Tel.: +54 11 4515 8200 | 8300
estimacionesagricolas@bc.org.ar
 Twitter: @estimacionesbc

ISSN 2408-4344t

We appreciate the contribution of our Network of Collaborators throughout the country.



RECENT ADVERSE EVENTS



METHODOLOGY

Findings of the survey conducted through January 12, 2021. The areas under analysis in this report account for 90% of the crops planted area. The regional development data is published once planting is complete. The national planting and harvest progress, as well as the phenological data of the crops derive from the final area projection, while the moisture and crop condition derive from the planting progress to

SOYBEAN 2021/22






| Regional participation in the total national area (%) | | | | | | | | | | | | | | | |
|---|------------------------|---------------|---|---------------------------------|---------|--------------------|-----------------|---------------|----------------|--------|-----------|-----|------|--------|-----|
| Regions | VII | VI | IX | III | II | IV | I | V | VIII | XII | X | XI | XIII | XIV | XV |
| Nat. Part. 2021/22 | 13,4 | 12,7 | 11,3 | 10,3 | 9,7 | 10,1 | 7,5 | 6,3 | 6,2 | 3,1 | 4,2 | 2,2 | 1,4 | 1,0 | 0,5 |
| Hist. Nat. Part. | 13,0 | 13,5 | 10,9 | 11,1 | 8,7 | 9,5 | 6,1 | 6,5 | 6,3 | 5,0 | 4,3 | 2,0 | 1,5 | 1,1 | 0,4 |
| Moisture condition, Crop condition & Development stages | | | | | | | | | | | | | | Week 3 | |
| Regions | Adverse event | Season | Moisture Condition Favourable / Optimum | Crop Condition Fair / Excellent | Planted | Leaves Development | Beginning bloom | Beginning pod | Beginning seed | Mature | Harvested | | | | |
| I - NWA | | 21/22 | 37,8 | 65,5 | 37,3 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 69,0 | 98,5 | 89,8 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 90,6 | - | - | - | - | - | - | - | - | - | - |
| II - NEA | Weeds | 21/22 | 44,3 | 76,3 | 84,8 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 53,7 | 95,0 | 82,0 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 87,8 | - | - | - | - | - | - | - | - | - | - |
| III - NC Cba | Pests - Dought | 21/22 | 11,0 | 77,9 | 100,0 | 93,0 | 42,2 | 12,5 | - | - | - | - | - | - | - |
| | | 20/21 | 59,9 | 79,6 | 100,0 | 97,0 | 27,4 | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 95,4 | 47,1 | 19,5 | 0,7 | - | - | - | - | - | - |
| IV - S Cba | Pests | 21/22 | 79,9 | 100,0 | 100,0 | 97,1 | 38,2 | 8,8 | - | - | - | - | - | - | - |
| | | 20/21 | 86,6 | 90,2 | 100,0 | 96,0 | 42,9 | 3,1 | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 95,2 | 52,4 | 19,6 | - | - | - | - | - | - | - |
| V - NC S Fe | Pests - Weeds - Dought | 21/22 | 16,4 | 66,4 | 88,1 | 81,4 | 40,8 | 4,4 | - | - | - | - | - | - | - |
| | | 20/21 | 51,5 | 70,4 | 100,0 | 95,6 | 40,7 | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,8 | 91,8 | 45,9 | 14,1 | 0,6 | - | - | - | - | - | - |
| VI - North Belt | Pests - Weeds - Dought | 21/22 | 37,4 | 64,5 | 100,0 | 97,0 | 46,8 | 22,5 | 1,2 | - | - | - | - | - | - |
| | | 20/21 | 68,0 | 74,3 | 100,0 | 98,3 | 58,7 | 7,3 | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 98,4 | 64,5 | 20,5 | - | - | - | - | - | - | - |
| VII - South Belt | Pests - Dought | 21/22 | 29,4 | 38,6 | 100,0 | 97,3 | 47,7 | 15,8 | - | - | - | - | - | - | - |
| | | 20/21 | 62,3 | 72,3 | 100,0 | 97,0 | 57,8 | 2,8 | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 96,4 | 64,7 | 15,3 | - | - | - | - | - | - | - |
| VIII - CE E. Rios | Pests - Dought | 21/22 | 1,8 | 37,8 | 97,7 | 95,6 | 34,0 | 4,2 | - | - | - | - | - | - | - |
| | | 20/21 | 38,9 | 69,9 | 100,0 | 98,0 | 42,9 | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,7 | 96,2 | 49,5 | 4,5 | - | - | - | - | - | - | - |
| IX - N LP - W BA | Pests | 21/22 | 75,2 | 93,6 | 98,4 | 92,0 | 65,7 | 6,1 | - | - | - | - | - | - | - |
| | | 20/21 | 77,2 | 71,3 | 100,0 | 92,4 | 51,7 | 2,2 | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 94,7 | 53,6 | 11,9 | - | - | - | - | - | - | - |
| XII - SE BA | Dought | 21/22 | 24,5 | 72,9 | 100,0 | 56,2 | 3,8 | - | - | - | - | - | - | - | - |
| | | 20/21 | 100,0 | 96,8 | 100,0 | 67,7 | 10,3 | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 71,8 | 13,5 | - | - | - | - | - | - | - | - |
| National | | 21/22 | 39,5 | 71,1 | 92,5 | 81,1 | 35,2 | 8,5 | 0,2 | - | - | - | - | - | - |
| | | 20/21 | 68,4 | 81,1 | 97,5 | 84,9 | 36,1 | 1,9 | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 98,4 | 86,8 | 44,1 | 11,8 | 0,1 | - | - | - | - | - | - |

FIRST SOYBEAN 2021/22

| Regional participation in the total national area (%) | | | | | | | | | | | | | | | |
|---|------|------|------|------|------|------|-----|------|-----|-----|-----|------|-----|-----|-----|
| Regions | VII | VI | III | IX | IV | II | I | VIII | V | X | XII | XIII | XIV | XI | XV |
| Nat. Part. 2021/22 | 13,6 | 12,1 | 11,7 | 11,8 | 11,4 | 10,3 | 8,4 | 6,7 | 6,1 | 2,8 | 1,4 | 2,1 | 0,6 | 0,6 | 0,5 |
| Hist. Nat. Part. | 13,9 | 13,3 | 11,5 | 11,4 | 10,2 | 9,3 | 6,2 | 7,2 | 6,2 | 3,5 | 3,4 | 2,0 | 0,9 | 0,7 | 0,4 |

| Moisture condition, Crop condition & Development stages | | | | | | | | | | | | Week 3 | | | |
|---|------------------------|---------------|---|---------------------------------|---------|--------------------|-----------------|---------------|----------------|--------|-----------|--------|--|--|--|
| Regions | Adverse event | Season | Moisture Condition Favourable / Optimum | Crop Condition Fair / Excellent | Planted | Leaves Development | Beginning bloom | Beginning pod | Beginning seed | Mature | Harvested | | | | |
| I - NWA | | 21/22 | 35,0 | - | 40,0 | - | - | - | - | - | - | | | | |
| | | 20/21 | 69,0 | - | 94,0 | - | - | - | - | - | - | | | | |
| | | 16/21 Average | - | - | 83,5 | - | - | - | - | - | - | | | | |
| II - NEA | Weeds | 21/22 | 40,0 | 85,0 | 75,0 | - | - | - | - | - | - | | | | |
| | | 20/21 | 52,0 | 95,0 | 86,0 | - | - | - | - | - | - | | | | |
| | | 16/21 Average | - | - | 85,3 | - | - | - | - | - | - | | | | |
| III - NC Cba | Pests - Dought | 21/22 | 6,0 | 80,0 | 100,0 | 98,0 | 54,0 | 17,0 | - | - | - | | | | |
| | | 20/21 | 56,0 | 74,0 | 100,0 | 100,0 | 35,0 | - | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 99,0 | 60,0 | 23,2 | 0,8 | - | - | | | | |
| IV - S Cba | Pests | 21/22 | 78,0 | 100,0 | 100,0 | 100,0 | 52,0 | 12,0 | - | - | - | | | | |
| | | 20/21 | 84,0 | 88,0 | 100,0 | 100,0 | 55,0 | 4,0 | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 64,6 | 24,4 | - | - | - | | | | |
| V - NC S Fe | Pests - Weeds - Dought | 21/22 | 20,0 | 70,0 | 90,0 | 90,0 | 62,0 | 7,0 | - | - | - | | | | |
| | | 20/21 | 42,0 | 62,0 | 100,0 | 100,0 | 64,0 | - | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 98,2 | 64,2 | 17,0 | 1,0 | - | - | | | | |
| VI - North Belt | Pests - Weeds - Dought | 21/22 | 40,0 | 60,0 | 100,0 | 100,0 | 75,0 | 36,0 | 2,0 | - | - | | | | |
| | | 20/21 | 64,0 | 71,0 | 100,0 | 100,0 | 83,0 | 11,0 | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 82,8 | 28,8 | - | - | - | | | | |
| VII - South Belt | Pests - Dought | 21/22 | 38,0 | 30,0 | 100,0 | 100,0 | 72,0 | 24,0 | - | - | - | | | | |
| | | 20/21 | 59,0 | 68,0 | 100,0 | 100,0 | 80,0 | 4,0 | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 80,4 | 19,0 | - | - | - | | | | |
| VIII - CE E. Rios | Pests - Dought | 21/22 | 1,0 | 37,0 | 100,0 | 100,0 | 48,0 | 6,0 | - | - | - | | | | |
| | | 20/21 | 30,0 | 64,0 | 100,0 | 100,0 | 57,0 | - | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 61,8 | 5,2 | - | - | - | | | | |
| IX - N LP - W BA | Pests | 21/22 | 74,0 | 93,0 | 100,0 | 100,0 | 95,0 | 9,0 | - | - | - | | | | |
| | | 20/21 | 73,0 | 64,0 | 100,0 | 100,0 | 72,0 | 3,0 | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 71,4 | 15,4 | - | - | - | | | | |
| National | | 21/22 | 39,6 | 70,7 | 93,0 | 86,9 | 53,2 | 13,0 | 0,2 | - | - | | | | |
| | | 20/21 | 63,1 | 76,6 | 98,0 | 90,6 | 51,5 | 2,7 | - | - | - | | | | |
| | | 16/21 Average | - | - | 97,9 | 92,7 | 58,9 | 15,7 | 0,2 | - | - | | | | |

CORN 2021/22



| Regional participation in the total national area (%) | | | | | | | | | | | | | | | |
|---|---------------|---------------|---|---------------------------------|---------|--------------------|-------------|---------|--------------|--------|-----------|-----|--------|-----|-----|
| Regions | III | IV | IX | II | VI | VII | I | X | VIII | XIII | XII | V | XI | XIV | XV |
| Nat. Part. 2019/20 | 13,7 | 11,8 | 10,3 | 9,9 | 9,0 | 6,9 | 5,8 | 4,9 | 4,7 | 4,5 | 3,6 | 3,6 | 2,4 | 1,7 | 0,5 |
| Hist. Nat. Part. | 14,5 | 12,4 | 10,7 | 11,0 | 10,0 | 7,9 | 6,4 | 5,1 | 4,6 | 4,9 | 3,6 | 3,8 | 2,7 | 1,8 | 0,5 |
| Moisture condition, Crop condition & Development stages | | | | | | | | | | | | | Week 3 | | |
| Regions | Adverse event | Season | Moisture Condition Favourable / Optimum | Crop Condition Fair / Excellent | Planted | Leaves Development | Pollination | Silking | Grain Growth | Mature | Harvested | | | | |
| I - NWA | | 21/22 | 65,0 | - | 29,0 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 95,0 | - | 57,0 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 56,0 | - | - | - | - | - | - | - | - | - | - |
| II - NEA | | 21/22 | 66,0 | - | 47,0 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 70,0 | - | 50,0 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 55,8 | - | - | - | - | - | - | - | - | - | - |
| III - NC Cba | Drought | 21/22 | 34,5 | - | 90,7 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 79,2 | - | 100,0 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 98,3 | - | - | - | - | - | - | - | - | - | - |
| IV - S Cba | | 21/22 | 69,6 | 91,2 | 100,0 | 70,6 | 34,7 | 6,2 | 0,4 | - | - | - | - | - | - |
| | | 20/21 | 88,8 | 91,4 | 100,0 | 81,5 | 12,0 | 11,8 | 2,2 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,2 | 79,1 | 24,0 | 20,2 | 3,1 | - | - | - | - | - | - |
| VI - North Belt | Drought | 21/22 | 24,1 | 36,7 | 100,0 | 95,5 | 89,5 | 84,8 | 28,9 | - | - | - | - | - | - |
| | | 20/21 | 65,6 | 70,4 | 100,0 | 99,3 | 85,3 | 85,0 | 38,3 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,9 | 97,0 | 87,7 | 86,4 | 45,3 | 1,2 | - | - | - | - | - |
| VII - South Belt | Drought | 21/22 | 39,4 | 57,7 | 100,0 | 95,9 | 89,6 | 85,6 | 18,1 | - | - | - | - | - | - |
| | | 20/21 | 61,4 | 60,2 | 100,0 | 98,2 | 86,1 | 86,0 | 25,8 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,8 | 97,0 | 86,0 | 85,0 | 26,1 | 0,2 | - | - | - | - | - |
| VIII - CE E. Rios | Drought | 21/22 | 3,4 | 18,5 | 88,8 | 69,0 | 44,2 | 38,0 | 18,6 | 1,9 | - | - | - | - | - |
| | | 20/21 | 53,0 | 72,2 | 98,1 | 65,9 | 38,0 | 38,0 | 22,8 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 95,4 | 78,7 | 57,7 | 57,6 | 34,2 | 4,0 | - | - | - | - | - |
| IX - N LP - W BA | | 21/22 | 70,8 | 82,4 | 98,7 | 85,7 | 55,7 | 30,9 | 1,1 | - | - | - | - | - | - |
| | | 20/21 | 84,0 | 78,8 | 100,0 | 89,8 | 40,0 | 40,0 | 4,0 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,6 | 90,8 | 52,8 | 50,2 | 6,3 | - | - | - | - | - | - |
| X - Central BA | | 21/22 | 63,0 | 77,6 | 100,0 | 82,6 | 52,6 | 30,0 | - | - | - | - | - | - | - |
| | | 20/21 | 95,6 | 92,2 | 100,0 | 91,6 | 40,0 | 36,0 | 2,0 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,8 | 87,6 | 46,3 | 40,6 | 4,6 | - | - | - | - | - | - |
| XII - SE BA | Drought | 21/22 | 17,2 | 54,8 | 100,0 | 76,9 | 41,1 | 12,2 | - | - | - | - | - | - | - |
| | | 20/21 | 100,0 | 97,5 | 100,0 | 92,5 | 34,0 | 16,0 | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,2 | 89,4 | 32,9 | 19,8 | - | - | - | - | - | - | - |
| XIII - San Luis | | 21/22 | 65,7 | 93,3 | 100,0 | 72,1 | 30,4 | 4,8 | 0,4 | - | - | - | - | - | - |
| | | 20/21 | 87,0 | 81,6 | 100,0 | 86,8 | 12,0 | 11,0 | 0,4 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,7 | 80,5 | 19,0 | 16,7 | 0,5 | - | - | - | - | - | - |
| National | | 21/22 | 47,2 | 63,9 | 86,4 | 67,3 | 42,4 | 26,4 | 6,1 | 0,2 | - | - | - | - | - |
| | | 20/21 | 79,7 | 83,9 | 90,9 | 74,8 | 31,0 | 29,3 | 8,9 | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 90,9 | 76,1 | 38,0 | 35,2 | 11,2 | 0,6 | - | - | - | - | - |

EARLY CORN 2021/22




| Regional participation in the total national area (%) | | | | | | | | | | | | | | | |
|---|------|------|------|------|-----|-----|------|-----|-----|-----|-----|-----|-----|------|-----|
| Regions | II | VI | VII | I | IX | X | VIII | XII | III | IV | V | XI | XIV | XIII | XV |
| Nat. Part. 2019/20 | 22,4 | 17,5 | 13,8 | 13,0 | 8,1 | 4,5 | 4,0 | 3,7 | 2,1 | 2,1 | 2,7 | 2,5 | 1,8 | 0,7 | 1,1 |
| Hist. Nat. Part. | 19,9 | 15,9 | 12,3 | 11,5 | 9,4 | 4,4 | 4,7 | 3,9 | 4,0 | 4,5 | 2,5 | 2,7 | 1,7 | 1,7 | 0,9 |

| Moisture condition, Crop condition & Development stages | | | | | | | | | | | | Week 3 | | | |
|---|---------------|---------------|---|---------------------------------|---------|--------------------|-------------|---------|--------------|--------|-----------|--------|--|--|--|
| Regions | Adverse event | Season | Moisture Condition Favourable / Optimum | Crop Condition Fair / Excellent | Planted | Leaves Development | Pollination | Silking | Grain Growth | Mature | Harvested | | | | |
| I - NWA | | 21/22 | 65,0 | - | 29,0 | - | - | - | - | - | - | | | | |
| | | 20/21 | 95,0 | - | 57,0 | - | - | - | - | - | - | | | | |
| | | 16/21 Average | - | - | 58,0 | - | - | - | - | - | - | | | | |
| II - NEA | | 21/22 | 66,0 | - | 47,0 | - | - | - | - | - | - | | | | |
| | | 20/21 | 70,0 | - | 50,0 | - | - | - | - | - | - | | | | |
| | | 16/21 Average | - | - | 56,6 | - | - | - | - | - | - | | | | |
| III - NC Cba | Drought | 21/22 | 41,0 | 54,0 | 100,0 | 100,0 | 98,0 | 81,0 | 9,0 | - | - | | | | |
| | | 20/21 | 72,0 | 65,0 | 100,0 | 100,0 | 100,0 | 89,0 | 19,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 98,6 | 90,8 | 17,2 | 0,4 | - | | | | |
| IV - S Cba | | 21/22 | 76,0 | 82,0 | 100,0 | 100,0 | 100,0 | 77,0 | 5,0 | - | - | | | | |
| | | 20/21 | 80,0 | 58,0 | 100,0 | 100,0 | 100,0 | 98,0 | 18,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 93,6 | 14,8 | - | - | | | | |
| VI - North Belt | Drought | 21/22 | 22,0 | 32,0 | 100,0 | 100,0 | 100,0 | 99,0 | 34,0 | - | - | | | | |
| | | 20/21 | 63,0 | 68,0 | 100,0 | 100,0 | 100,0 | 100,0 | 45,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 99,0 | 51,6 | 1,4 | - | | | | |
| VII - South Belt | Drought | 21/22 | 39,0 | 57,0 | 100,0 | 100,0 | 100,0 | 99,0 | 21,0 | - | - | | | | |
| | | 20/21 | 58,0 | 55,0 | 100,0 | 100,0 | 100,0 | 100,0 | 30,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 99,0 | 30,2 | 0,2 | - | | | | |
| VIII - CE E. Rios | Drought | 21/22 | - | 3,0 | 100,0 | 100,0 | 100,0 | 100,0 | 49,0 | 5,0 | - | | | | |
| | | 20/21 | 23,0 | 36,0 | 100,0 | 100,0 | 100,0 | 100,0 | 60,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 59,4 | 6,4 | - | | | | |
| IX - N LP - W BA | | 21/22 | 76,0 | 85,0 | 100,0 | 100,0 | 96,0 | 79,0 | 3,0 | - | - | | | | |
| | | 20/21 | 75,0 | 56,0 | 100,0 | 100,0 | 100,0 | 100,0 | 10,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 97,0 | 94,8 | 11,8 | - | - | | | | |
| X - Central BA | | 21/22 | 69,0 | 68,0 | 100,0 | 100,0 | 94,0 | 75,0 | - | - | - | | | | |
| | | 20/21 | 89,0 | 88,0 | 100,0 | 100,0 | 100,0 | 90,0 | 5,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 95,0 | 84,8 | 9,6 | - | - | | | | |
| XII - SE BA | Drought | 21/22 | 10,0 | 63,0 | 100,0 | 100,0 | 79,0 | 27,0 | - | - | - | | | | |
| | | 20/21 | 100,0 | 100,0 | 100,0 | 100,0 | 68,0 | 32,0 | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 54,0 | 32,4 | - | - | - | | | | |
| National | | 21/22 | 42,6 | 54,4 | 78,3 | 66,5 | 59,0 | 51,5 | 12,0 | 0,4 | - | | | | |
| | | 20/21 | 71,5 | 72,1 | 82,7 | 71,9 | 61,8 | 58,6 | 17,7 | - | - | | | | |
| | | 16/21 Average | - | - | 86,7 | 79,4 | 66,2 | 62,2 | 19,7 | 1,0 | - | | | | |

LATE CORN 2021/22

| Regional participation in the total national area (%) | | | | | | | | | | | | | | | |
|---|---------------|---------------|---|---------------------------------|---------|--------------------|-------------|---------|--------------|--------|-----------|-----|-----|--------|-----|
| Regions | III | IV | IX | XIII | X | VIII | V | XII | VI | XI | VII | XIV | II | I | XV |
| Nat. Part. 2021/22 | 25,7 | 21,9 | 13,5 | 8,5 | 6,1 | 5,9 | 4,8 | 4,1 | 2,8 | 2,7 | 2,0 | 2,0 | 0,0 | 0,0 | 0,0 |
| Hist. Nat. Part. | 27,6 | 22,1 | 12,3 | 8,8 | 6,0 | 4,5 | 5,4 | 3,3 | 2,8 | 2,6 | 2,4 | 2,0 | 0,0 | 0,0 | 0,0 |
| Moisture condition, Crop condition & Development stages | | | | | | | | | | | | | | Week 3 | |
| Regions | Adverse event | Season | Moisture Condition Favourable / Optimum | Crop Condition Fair / Excellent | Planted | Leaves Development | Pollination | Silking | Grain Growth | Mature | Harvested | | | | |
| III - NC Cba | Drought | 21/22 | 34,0 | - | 90,0 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 80,0 | - | 100,0 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 98,6 | - | - | - | - | - | - | - | - | - | - |
| IV - S Cba | | 21/22 | 69,0 | 92,0 | 100,0 | 68,0 | 29,0 | - | - | - | - | - | - | - | - |
| | | 20/21 | 90,0 | 96,0 | 100,0 | 79,0 | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 99,6 | 73,2 | 3,2 | 0,2 | - | - | - | - | - | - | - |
| V - NC S Fe | Drought | 21/22 | 15,0 | 53,0 | 53,0 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 75,0 | 88,0 | 85,0 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 77,4 | - | - | - | - | - | - | - | - | - | - |
| VIII - EC E. Rios | Drought | 21/22 | 6,0 | 30,0 | 82,0 | - | - | - | - | - | - | - | - | - | - |
| | | 20/21 | 72,0 | 95,0 | 97,0 | - | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 93,6 | - | - | - | - | - | - | - | - | - | - |
| IX - N LP - W BA | | 21/22 | 68,0 | 81,0 | 98,0 | 78,0 | 34,0 | 5,0 | - | - | - | - | - | - | - |
| | | 20/21 | 90,0 | 94,0 | 100,0 | 83,0 | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 80,6 | 3,4 | 0,2 | - | - | - | - | - | - | - |
| X - Central BA | | 21/22 | 59,0 | 84,0 | 100,0 | 71,0 | 25,0 | - | - | - | - | - | - | - | - |
| | | 20/21 | 100,0 | 95,0 | 100,0 | 86,0 | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 77,0 | 1,4 | - | - | - | - | - | - | - | - |
| XII - SE BA | Drought | 21/22 | 23,0 | 48,0 | 100,0 | 58,0 | 10,0 | - | - | - | - | - | - | - | - |
| | | 20/21 | 100,0 | 95,0 | 100,0 | 85,0 | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 71,0 | 0,4 | - | - | - | - | - | - | - | - |
| XIII - San Luis | | 21/22 | 68,0 | 95,0 | 100,0 | 70,0 | 26,0 | - | - | - | - | - | - | - | - |
| | | 20/21 | 88,0 | 84,0 | 100,0 | 85,0 | - | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 100,0 | 75,6 | 1,6 | 0,2 | - | - | - | - | - | - | - |
| National | | 21/22 | 50,6 | 71,1 | 93,7 | 59,4 | 22,3 | 0,8 | - | - | - | - | - | - | - |
| | | 20/21 | 86,5 | 93,8 | 99,1 | 77,8 | 0,1 | - | - | - | - | - | - | - | - |
| | | 16/21 Average | - | - | 97,4 | 71,5 | 1,8 | 0,2 | - | - | - | - | - | - | - |

SUNFLOWER 2021/22



| Regional participation in the total national area (%) | | | | | | | | | | | | | | | |
|---|---------------|---------------|---|---------------------------------|---------|--------------------|-----------------|-----------------|------------------|--------|-----------|------|-----|--------|-----|
| Regions | XII | XI | II | V | IX | XIV | X | IV | XIII | VI | VII | VIII | XV | III | I |
| Nat. Part. 2019/20 | 23,0 | 21,6 | 10,6 | 14,2 | 12,4 | 7,3 | 4,5 | 1,7 | 1,4 | 0,9 | 0,8 | 0,7 | 0,5 | 0,4 | 0,0 |
| Hist. Nat. Part. | 23,3 | 22,3 | 17,5 | 14,1 | 7,7 | 6,1 | 3,9 | 1,3 | 1,4 | 0,7 | 0,5 | 0,4 | 0,5 | 0,3 | 0,0 |
| Moisture condition, Crop condition & Development stages | | | | | | | | | | | | | | Week 3 | |
| Regions | Adverse event | Season | Moisture Condition Favourable / Optimum | Crop Condition Fair / Excellent | Planted | Leaves Development | Flowerbud Stage | Flowering stage | Seed development | Mature | Harvested | | | | |
| II - NEA | | 21/22 | 52,0 | 92,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 60,0 | | | |
| | | 20/21 | 9,0 | 2,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 87,5 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 73,2 | | | |
| V - NC S Fe | | 21/22 | 60,0 | 80,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 93,0 | 30,0 | | | | |
| | | 20/21 | 66,0 | 86,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 88,0 | 10,0 | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 91,2 | 28,5 | | | | |
| IX - N LP - W BA | | 21/22 | 83,0 | 100,0 | 100,0 | 100,0 | 100,0 | 72,0 | 30,0 | - | - | | | | |
| | | 20/21 | 80,0 | 85,0 | 100,0 | 100,0 | 100,0 | 71,0 | 28,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 68,2 | 31,2 | 0,4 | - | - | | | |
| X - Central BA | Drought | 21/22 | 49,0 | 97,0 | 100,0 | 100,0 | 96,0 | 68,0 | 28,0 | - | - | | | | |
| | | 20/21 | 90,0 | 99,0 | 100,0 | 100,0 | 100,0 | 65,0 | 22,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 95,6 | 64,0 | 25,0 | - | - | | | | |
| XI - SW BA - S LP | | 21/22 | 67,0 | 100,0 | 100,0 | 100,0 | 84,0 | 47,0 | 10,0 | - | - | | | | |
| | | 20/21 | 95,0 | 93,0 | 100,0 | 100,0 | 84,0 | 46,0 | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 81,2 | 46,4 | 5,8 | - | - | | | | |
| XII - SE BA | Drought | 21/22 | 7,0 | 60,0 | 100,0 | 100,0 | 86,0 | 50,0 | 9,0 | - | - | | | | |
| | | 20/21 | 88,0 | 94,0 | 100,0 | 100,0 | 89,0 | 50,0 | - | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 82,8 | 55,4 | 7,4 | - | - | | | | |
| XIV - Cuenca Sal. | Drought | 21/22 | - | 40,0 | 100,0 | 100,0 | 100,0 | 78,0 | 25,0 | - | - | | | | |
| | | 20/21 | 98,0 | 95,0 | 100,0 | 100,0 | 100,0 | 65,0 | 19,0 | - | - | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 96,6 | 63,6 | 27,8 | - | - | | | | |
| National | | 21/22 | 43,2 | 80,7 | 100,0 | 100,0 | 93,1 | 70,0 | 39,3 | 24,8 | 10,4 | | | | |
| | | 20/21 | 86,6 | 91,4 | 100,0 | 100,0 | 93,0 | 63,7 | 25,5 | 16,5 | 5,0 | | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 91,4 | 70,6 | 41,9 | 31,1 | 16,2 | | | | |

WHEAT 2021/22

| Regional participation in the total national area (%) | | | | | | | | | | | | | | | |
|---|---------------|---------------|---|---------------------------------|---------|-----------|----------------|---------|----------|--------|-----------|--------|-----|-----|------|
| Regions | VI | III | XII | VII | XI | IX | V | IV | I | II | X | VIII | XIV | XV | XIII |
| Nat. Part. 2021/22 | 12,9 | 10,0 | 9,9 | 9,8 | 8,8 | 8,0 | 7,4 | 6,8 | 6,6 | 6,6 | 5,5 | 5,1 | 2,1 | 0,4 | 0,2 |
| Hist. Nat. Part. | 13,5 | 10,5 | 9,8 | 9,8 | 8,8 | 8,3 | 7,2 | 6,7 | 6,2 | 6,7 | 5,5 | 4,8 | 1,8 | 0,3 | 0,2 |
| Moisture condition, Crop condition & Development stages | | | | | | | | | | | | Week 3 | | | |
| Regions | Adverse event | Season | Moisture Condition Favourable / Optimum | Crop Condition Fair / Excellent | Planted | Tillering | Stem Extension | Heading | Ripening | Mature | Harvested | | | | |
| I - NWA | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| II - NEA | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| III - NC Cba | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| IV - S Cba | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| V - NC S Fe | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| VI - North Belt | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| VII - South Belt | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| IX - N LP - W BA | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| X - Central BA | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| XI - SW BA - S LP | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| XII - SE BA | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| National | | 21/22 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 20/21 | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |
| | | 16/21 Average | - | - | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | 100,0 | | | |

Wheat 2021/22 Development in Argentina

Crop Condition: ■ Poor/Very poor ■ Fair ■ Excellent/Good
 Development Stages: — Planting — Leaves Development — Tillering — Steam Extension — Ripening — Mature - - - Harvest

