

CROP CONDITION & DEVELOPMENT STAGES


JANUARY, 9
2020



BUENOS AIRES GRAIN EXCHANGE
DEPARTMENT OF AGRICULTURAL
ESTIMATES

ISSN 2591-4715

DEPARTMENT & REGIONS



- I - NOA (Noroeste Argentino)
- II - NEA (Noreste Argentino)
- III - Centro-Norte de Córdoba
- IV - Sur de Córdoba
- V - Centro-Norte de Santa Fe
- VI - Núcleo Norte
- VII - Núcleo Sur
- VIII - Centro-Este de Entre Ríos
- IX - Norte de La Pampa - Oeste de Buenos Aires
- X - Centro de Buenos Aires
- XI - Sudoeste de Buenos Aires - Sur de La Pampa
- XII - Sudeste de Buenos Aires
- XIII - San Luis
- XIV - Cuenca del Salado
- XV - Corrientes - Misiones

HEAD OF DEPARTMENT

Esteban J Copati
ecopati@bc.org.ar

CROP ANALYST

Martin López
martinlopez@bc.org.ar

CROP ANALYST

Daniela A. Venturino
dventurino@bc.org.ar

CROP CONDITION ANALYST

Sofía Console Insúa
sconsoleinsua@bc.org.ar
Soybean, Wheat, Sunflower

Matías Mihura
mmihura@bc.org.ar
Corn

CONTACT

123 Corrientes Avenue
C1043AAB - CABA
Phone: +54 11 4515 8200 | 8300
estimacionesagricolas@bc.org.ar
Twitter: @estimacionesbc

ISSN 2591-4715

CROP CONDITION & DEVELOPMENT STAGES

Findings of the survey conducted through January, 8 2020. The areas under analysis in this report account for 90% of the crops planted area. The regional development data is published once planting is complete. The national planting and harvest progress, as well as the phenological data of the crops derive from the final area projection, while the moisture and crop condition derive from the planting progress to date.

First Soybean

Season 2019/2020

First Soybean Planted: as of January 8, 2020

Regions	January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I NWA	86.0	24.7	45.8	74.8
II NEA	74.0	45.4	79.3	70.4
III NCnt Cba	99.0	97.0	99.4	99.4
IV S Cba	100.0	96.8	100.0	99.8
V NCnt SFe	98.0	94.8	97.5	97.0
VI North Belt	100.0	100.0	100.0	99.8
VII South Belt	99.0	97.4	100.0	99.8
VIII ECnt ER	98.0	96.8	100.0	99.4
IX N LP-W BA	99.0	95.1	100.0	99.2
X Cnt BA	96.0	78.5	91.3	98.4
XI SW BA-S LP	97.0	60.4	79.5	91.6
XII SE BA	94.0	52.2	83.2	94.0
XIII SL	99.0	99.3	100.0	99.8
XIV Cuenca Sal	79.0	76.0	83.2	95.0
XV Others	94.0	96.0	97.1	98.4
National	96.0	84.3	93.2	95.4

First Soybean Condition: as of January 8, 2020

Regions	Very Poor %	Poor %	Fair %	Good %	Excellent %
I NWA	-	0.3	40.8	41.9	17.1
II NEA	-	-	34.8	43.1	22.1
III NCnt Cba	-	1.4	38.1	56.9	3.6
IV S Cba	-	0.7	43.1	52.7	3.4
V NCnt SFe	-	-	13.5	64.6	21.9
VI North Belt	-	-	30.0	53.0	17.1
VII South Belt	-	-	18.7	72.3	9.0
VIII ECnt ER	-	9.0	53.8	29.3	8.0
IX N LP-W BA	-	0.7	52.8	46.0	0.6
X Cnt BA	-	5.6	76.9	17.5	-
XI SW BA-S LP	0.8	7.6	39.5	52.2	-
XII SE BA	-	-	36.5	58.8	4.7
XIII SL	-	2.8	81.0	16.2	-
XIV Cuenca Sal	-	5.6	64.5	29.9	-
XV Others	-	-	24.9	68.5	6.6
National	0.0	1.5	38.0	51.4	9.1
Nat. Jan 9, 2019	0.8	8.3	42.1	42.2	6.6

CROP CONDITION & DEVELOPMENT STAGES

First Soybean Moisture Condition: as of January 8, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	5.8	43.4	50.9	-
II	NEA	-	-	22.9	76.3	0.7
III	NCnt Cba	-	3.4	65.7	30.9	-
IV	S Cba	-	14.7	73.4	11.9	-
V	NCnt SFe	-	2.5	49.9	40.5	7.1
VI	North Belt	-	3.1	49.5	46.7	0.7
VII	South Belt	-	2.5	70.2	27.3	-
VIII	ECnt ER	-	-	50.3	48.0	1.8
IX	N LP-W BA	-	9.0	74.2	16.9	-
X	Cnt BA	-	24.4	65.0	10.6	-
XI	SW BA-S LP	3.0	26.4	50.6	20.1	-
XII	SE BA	1.2	12.0	56.8	30.0	-
XIII	SL	4.5	25.4	57.7	12.3	-
XIV	Cuenca Sal	-	18.7	59.4	21.9	-
XV	Others	-	-	34.5	61.8	3.7
National		0.2	6.7	58.7	33.7	0.8
Nat. Jan 9, 2019		-	3.2	45.7	37.4	13.8

First Soybean Leaf Development: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	85.0	50.0	76.0	89.6
IV	S Cba	95.0	98.0	100.0	95.0
V	NCnt SFe	85.0	100.0	100.0	85.6
VI	North Belt	100.0	100.0	100.0	92.4
VII	South Belt	100.0	100.0	100.0	99.0
VIII	ECnt ER	96.0	100.0	100.0	98.2
IX	N LP-W BA	99.0	100.0	100.0	99.2
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	82.0	100.0	100.0	86.4
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		87.4	86.3	87.7	87.1

CROP CONDITION & DEVELOPMENT STAGES

First Soybean Beginning Bloom: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	46.0	36.0	48.0	61.4
IV	S Cba	50.0	39.0	58.0	56.6
V	NCnt SFe	38.0	40.0	50.0	59.6
VI	North Belt	63.0	50.0	68.0	66.4
VII	South Belt	55.0	50.0	70.0	66.2
VIII	ECnt ER	38.0	37.0	47.0	60.0
IX	N LP-W BA	50.0	38.0	56.0	50.6
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	16.0	-	10.0	16.7
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		40.0	32.9	46.4	49.8

First Soybean Beginning Pod: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	11.0	-	-	19.1
IV	S Cba	8.4	-	13.3	16.3
V	NCnt SFe	-	0.6	-	14.7
VI	North Belt	13.0	-	7.8	13.8
VII	South Belt	6.8	-	4.4	11.0
VIII	ECnt ER	-	-	-	7.3
IX	N LP-W BA	4.5	-	7.2	5.3
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		5.2	0.0	4.8	9.1

CROP CONDITION & DEVELOPMENT STAGES

Second Soybean

Second Soybean Planted: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	75.0	10.0	38.0	68.6
II	NEA	68.0	40.7	80.6	67.6
III	NCnt Cba	98.0	90.3	98.0	97.6
IV	S Cba	100.0	87.6	100.0	98.4
V	NCnt SFe	94.0	86.0	93.2	92.2
VI	North Belt	99.0	100.0	100.0	98.8
VII	South Belt	96.0	90.0	100.0	98.6
VIII	ECnt ER	90.0	87.2	100.0	96.0
IX	N LP-W BA	94.0	82.4	100.0	96.4
X	Cnt BA	92.0	60.0	83.9	95.0
XI	SW BA-S LP	95.0	45.0	70.0	88.8
XII	SE BA	90.0	18.3	67.0	87.6
XIII	SL	87.0	86.6	100.0	97.0
XIV	Cuenca Sal	68.0	50.0	65.0	89.6
XV	Others	88.0	86.0	90.0	95.6
National		92.0	27.6	30.8	90.8

Second Soybean Condition: as of January 8, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	1.0	97.0	2.0	-
II	NEA	-	-	93.0	7.0	-
III	NCnt Cba	-	-	8.0	80.0	12.0
IV	S Cba	-	-	4.0	83.0	13.0
V	NCnt SFe	-	-	-	62.0	38.0
VI	North Belt	-	-	-	66.0	34.0
VII	South Belt	-	-	4.0	76.0	20.0
VIII	ECnt ER	-	-	8.0	60.0	32.0
IX	N LP-W BA	-	-	5.0	93.0	2.0
X	Cnt BA	-	-	85.0	15.0	-
XI	SW BA-S LP	-	4.0	10.0	86.0	-
XII	SE BA	-	-	53.0	47.0	-
XIII	SL	-	-	62.0	38.0	-
XIV	Cuenca Sal	-	-	47.0	53.0	-
XV	Others	-	-	41.0	59.0	-
National		-	0.2	20.8	63.3	15.8
Nat. Jan 9, 2019		0.8	9.6	49.2	37.1	3.4

CROP CONDITION & DEVELOPMENT STAGES

Second Soybean Moisture Condition: as of January 8, 2020

Regions	Dry %	Poor %	Favourable %	Optimum %	Wet %
I NWA	-	20.0	64.0	16.0	-
II NEA	-	-	45.0	55.0	-
III NCnt Cba	-	2.0	65.0	33.0	-
IV S Cba	-	25.0	69.0	6.0	-
V NCnt SFe	-	7.0	91.0	2.0	-
VI North Belt	-	9.0	58.0	33.0	-
VII South Belt	-	4.0	79.0	17.0	-
VIII ECnt ER	-	-	87.0	12.0	1.0
IX N LP-W BA	-	22.0	72.0	6.0	-
X Cnt BA	-	29.0	66.0	5.0	-
XI SW BA-S LP	1.0	16.0	51.0	32.0	-
XII SE BA	-	2.0	55.0	43.0	-
XIII SL	14.0	52.0	34.0	-	-
XIV Cuenca Sal	-	-	50.0	50.0	-
XV Others	-	-	47.0	53.0	-
National	0.1	10.5	67.4	22.0	0.1
Nat. Jan 9, 2019	-	4.2	46.6	35.4	13.8

Second Soybean Leaf Development: as of January 8, 2020

Regions	January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I NWA	-	-	-	-
II NEA	-	-	-	-
III NCnt Cba	-	-	-	-
IV S Cba	64.0	31.0	66.0	57.4
V NCnt SFe	-	-	-	-
VI North Belt	81.0	34.0	73.0	77.0
VII South Belt	65.0	33.0	50.0	56.8
VIII ECnt ER	64.0	30.0	60.0	65.2
IX N LP-W BA	71.0	27.0	50.0	61.6
X Cnt BA	-	-	-	-
XI SW BA-S LP	-	-	-	-
XII SE BA	-	-	-	-
XIII SL	-	-	-	-
XIV Cuenca Sal	-	-	-	-
XV Others	-	-	-	-
National	61.0	22.9	47.8	49.4

CROP CONDITION & DEVELOPMENT STAGES

Soybean

Soybean Planted: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	86.0	24.7	45.8	74.8
II	NEA	74.0	45.4	79.3	70.4
III	NCnt Cba	99.0	97.0	99.4	99.4
IV	S Cba	100.0	96.8	100.0	99.8
V	NCnt SFe	98.0	94.8	97.5	97.0
VI	North Belt	100.0	100.0	100.0	99.8
VII	South Belt	99.0	97.4	100.0	99.8
VIII	ECnt ER	98.0	96.8	100.0	99.4
IX	N LP-W BA	99.0	95.1	100.0	99.2
X	Cnt BA	96.0	78.5	91.3	98.4
XI	SW BA-S LP	97.0	60.4	79.5	91.6
XII	SE BA	94.0	52.2	83.2	94.0
XIII	SL	99.0	99.3	100.0	99.8
XIV	Cuenca Sal	79.0	76.0	83.2	95.0
XV	Others	94.0	96.0	97.1	98.4
National		96.0	84.3	93.2	95.4

Soybean Condition: as of January 8, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	0.3	40.8	41.9	17.1
II	NEA	-	-	34.8	43.1	22.1
III	NCnt Cba	-	1.4	38.1	56.9	3.6
IV	S Cba	-	0.7	43.1	52.7	3.4
V	NCnt SFe	-	-	13.5	64.6	21.9
VI	North Belt	-	-	30.0	53.0	17.1
VII	South Belt	-	-	18.7	72.3	9.0
VIII	ECnt ER	-	9.0	53.8	29.3	8.0
IX	N LP-W BA	-	0.7	52.8	46.0	0.6
X	Cnt BA	-	5.6	76.9	17.5	-
XI	SW BA-S LP	0.8	7.6	39.5	52.2	-
XII	SE BA	-	-	36.5	58.8	4.7
XIII	SL	-	2.8	81.0	16.2	-
XIV	Cuenca Sal	-	5.6	64.5	29.9	-
XV	Others	-	-	24.9	68.5	6.6
National		0.0	1.5	38.0	51.4	9.1
Nat. Jan 9, 2019		0.8	8.3	42.1	42.2	6.6

CROP CONDITION & DEVELOPMENT STAGES

Soybean Moisture Condition: as of January 8, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	5.8	43.4	50.9	-
II	NEA	-	-	22.9	76.3	0.7
III	NCnt Cba	-	3.4	65.7	30.9	-
IV	S Cba	-	14.7	73.4	11.9	-
V	NCnt SFe	-	2.5	49.9	40.5	7.1
VI	North Belt	-	3.1	49.5	46.7	0.7
VII	South Belt	-	2.5	70.2	27.3	-
VIII	ECnt ER	-	-	50.3	48.0	1.8
IX	N LP-W BA	-	9.0	74.2	16.9	-
X	Cnt BA	-	24.4	65.0	10.6	-
XI	SW BA-S LP	3.0	26.4	50.6	20.1	-
XII	SE BA	1.2	12.0	56.8	30.0	-
XIII	SL	4.5	25.4	57.7	12.3	-
XIV	Cuenca Sal	-	18.7	59.4	21.9	-
XV	Others	-	-	34.5	61.8	3.7
National		0.2	6.7	58.7	33.7	0.8
Nat. Jan 9, 2019		-	3.2	45.7	37.4	13.8

Soybean Leaf Development: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	-	-	-	-
IV	S Cba	87.6	72.3	73.8	89.4
V	NCnt SFe	-	-	-	-
VI	North Belt	94.0	65.1	65.1	89.6
VII	South Belt	91.4	73.5	73.5	92.8
VIII	ECnt ER	89.3	75.0	75.0	93.2
IX	N LP-W BA	92.2	72.4	72.4	91.7
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		79.0	58.3	59.3	79.0

CROP CONDITION & DEVELOPMENT STAGES

Early Corn

Early Corn Planted: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	55.0	21.0	30.0	41.8
II	NEA	38.0	23.0	37.0	44.4
III	NCnt Cba	100.0	100.0	100.0	100.0
IV	S Cba	100.0	100.0	100.0	100.0
V	NCnt SFe	100.0	100.0	100.0	100.0
VI	North Belt	100.0	100.0	100.0	100.0
VII	South Belt	100.0	100.0	100.0	100.0
VIII	ECnt ER	100.0	100.0	100.0	100.0
IX	N LP-W BA	100.0	100.0	100.0	100.0
X	Cnt BA	100.0	100.0	100.0	100.0
XI	SW BA-S LP	100.0	100.0	100.0	100.0
XII	SE BA	100.0	100.0	100.0	100.0
XIII	SL	100.0	100.0	100.0	100.0
XIV	Cuenca Sal	100.0	100.0	100.0	100.0
XV	Others	100.0	30.0	35.0	100.0
National		82.0	80.0	83.1	88.0

Early Corn Condition: as of January 8, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	3.3	55.0	41.7	-
II	NEA	-	-	62.5	37.5	-
III	NCnt Cba	-	3.3	65.0	28.3	3.3
IV	S Cba	-	9.2	52.5	37.5	0.8
V	NCnt SFe	-	1.1	10.0	78.9	10.0
VI	North Belt	-	2.0	22.0	67.0	9.0
VII	South Belt	-	7.8	44.4	34.4	13.3
VIII	ECnt ER	-	-	60.0	37.1	2.9
IX	N LP-W BA	-	6.4	15.7	72.9	5.0
X	Cnt BA	-	6.3	23.8	68.8	1.3
XI	SW BA-S LP	5.0	44.4	35.0	13.3	2.2
XII	SE BA	-	8.3	71.7	20.0	-
XIII	SL	-	15.0	55.0	30.0	-
XIV	Cuenca Sal	-	15.0	30.0	55.0	-
XV	Others	-	10.0	50.0	40.0	-
National		0.2	6.3	39.3	48.8	5.4
Nat. Jan 9, 2019		0.3	4.9	36.1	47.7	11.0

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Moisture Condition: as of January 8, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	20.0	50.0	30.0	-
II	NEA	-	2.5	45.0	52.5	-
III	NCnt Cba	-	8.3	50.0	41.7	-
IV	S Cba	-	8.3	23.3	68.3	-
V	NCnt SFe	-	14.4	71.1	14.4	-
VI	North Belt	-	15.0	43.0	41.0	1.0
VII	South Belt	-	6.7	38.9	53.3	1.1
VIII	ECnt ER	-	-	27.1	72.9	-
IX	N LP-W BA	-	2.9	14.3	82.9	-
X	Cnt BA	-	8.8	35.0	56.3	-
XI	SW BA-S LP	15.6	61.1	15.6	7.8	-
XII	SE BA	8.3	26.7	40.0	25.0	-
XIII	SL	-	-	80.0	20.0	-
XIV	Cuenca Sal	10.0	15.0	15.0	60.0	-
XV	Others	-	-	50.0	50.0	-
National		1.1	10.7	36.9	50.9	0.4
Nat. Jan 9, 2019		0.0	2.8	45.8	44.9	6.4

Early Corn Leaf Development: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	33.0	9.0	14.0	17.2
II	NEA	24.0	10.0	15.0	13.2
III	NCnt Cba	100.0	100.0	100.0	86.6
IV	S Cba	100.0	100.0	100.0	94.6
V	NCnt SFe	100.0	100.0	100.0	91.4
VI	North Belt	100.0	100.0	100.0	97.4
VII	South Belt	100.0	100.0	100.0	97.6
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100.0	100.0	100.0	98.8
X	Cnt BA	100.0	100.0	100.0	99.6
XI	SW BA-S LP	-	-	-	-
XII	SE BA	100.0	100.0	100.0	98.6
XIII	SL	100.0	100.0	100.0	97.0
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		79.2	76.8	78.1	72.8

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Pollination: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	0.4
II	NEA	-	-	-	0.4
III	NCnt Cba	100.0	85.0	97.0	80.2
IV	S Cba	100.0	82.0	100.0	88.0
V	NCnt SFe	100.0	100.0	100.0	91.4
VI	North Belt	100.0	99.0	100.0	96.6
VII	South Belt	100.0	96.0	100.0	96.6
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	98.0	82.0	91.0	86.0
X	Cnt BA	93.0	72.0	89.0	85.4
XI	SW BA-S LP	-	-	-	-
XII	SE BA	30.0	20.0	25.0	29.2
XIII	SL	80.0	40.0	65.0	72.8
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		66.9	61.7	67.8	60.0

Early Corn Silking: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	95.0	70.0	84.0	72.2
IV	S Cba	97.0	60.0	95.0	79.0
V	NCnt SFe	100.0	99.0	100.0	91.4
VI	North Belt	100.0	98.0	100.0	91.4
VII	South Belt	98.0	86.0	96.0	92.0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	93.0	64.0	83.0	82.2
X	Cnt BA	70.0	49.0	57.0	65.0
XI	SW BA-S LP	-	-	-	-
XII	SE BA	15.0	14.0	16.0	13.4
XIII	SL	65.0	29.0	52.0	49.0
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		63.2	53.8	61.7	53.8

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Grain Growth: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	18.0	4.0	12.0	9.4
IV	S Cba	5.0	1.0	6.0	5.8
V	NCnt SFe	58.0	51.0	57.0	56.6
VI	North Belt	26.0	14.0	30.0	27.2
VII	South Belt	15.0	7.0	14.0	16.6
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	2.0	-	5.0	3.8
X	Cnt BA	1.0	-	1.0	1.0
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	4.0	-	1.0	0.8
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		12.5	6.4	11.9	10.6

Late Corn

Late Corn Planted: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	95.0	92.0	100.0	91.2
IV	S Cba	97.0	96.0	100.0	94.4
V	NCnt SFe	40.0	40.0	56.0	57.6
VI	North Belt	96.0	100.0	100.0	97.2
VII	South Belt	92.0	100.0	100.0	96.4
VIII	ECnt ER	60.0	61.0	80.0	82.2
IX	N LP-W BA	96.0	92.0	100.0	97.4
X	Cnt BA	98.0	93.0	100.0	98.8
XI	SW BA-S LP	95.0	98.0	100.0	99.0
XII	SE BA	92.0	98.0	100.0	97.8
XIII	SL	98.0	92.0	100.0	95.2
XIV	Cuenca Sal	95.0	91.0	100.0	98.4
XV	Others	-	-	-	-
National		91.0	89.3	96.7	84.8

CROP CONDITION & DEVELOPMENT STAGES

Late Corn Condition: as of January 8, 2020

Regions	Very Poor %	Poor %	Fair %	Good %	Excellent %
I NWA	-	-	-	-	-
II NEA	-	-	-	-	-
III NCnt Cba	-	3.0	49.0	41.0	7.0
IV S Cba	-	1.0	28.0	58.0	13.0
V NCnt SFe	-	-	13.0	66.0	21.0
VI North Belt	-	1.0	15.0	73.0	11.0
VII South Belt	-	1.0	74.0	19.0	6.0
VIII ECnt ER	-	-	60.0	40.0	-
IX N LP-W BA	-	1.0	19.0	61.0	19.0
X Cnt BA	-	5.0	20.0	69.0	6.0
XI SW BA-S LP	5.0	14.0	44.0	37.0	-
XII SE BA	-	5.0	72.0	23.0	-
XIII SL	-	14.0	66.0	20.0	-
XIV Cuenca Sal	5.0	20.0	25.0	33.0	17.0
XV Others	-	-	-	-	-
National	0.2	3.9	40.2	46.7	8.9
Nat. Jan 9, 2019	0.6	6.2	50.5	37.8	5.0

Late Corn Moisture Condition: as of January 8, 2020

Regions	Dry %	Poor %	Favourable %	Optimum %	Wet %
I NWA	-	-	-	-	-
II NEA	-	-	-	-	-
III NCnt Cba	-	2.0	51.0	47.0	-
IV S Cba	-	5.0	30.0	65.0	-
V NCnt SFe	-	5.0	60.0	31.0	4.0
VI North Belt	-	20.0	35.0	45.0	-
VII South Belt	-	20.0	38.0	40.0	2.0
VIII ECnt ER	-	2.0	43.0	54.0	1.0
IX N LP-W BA	-	3.0	37.0	60.0	-
X Cnt BA	1.0	5.0	49.0	45.0	-
XI SW BA-S LP	14.0	50.0	36.0	-	-
XII SE BA	1.0	35.0	29.0	35.0	-
XIII SL	1.0	45.0	44.0	10.0	-
XIV Cuenca Sal	2.0	23.0	45.0	30.0	-
XV Others	-	-	-	-	-
National	0.6	11.1	41.5	46.6	0.2
Nat. Jan 9, 2019	0.0	16.2	53.8	23.7	6.2

CROP CONDITION & DEVELOPMENT STAGES

Late Corn Leaf Development: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	70.0	33.3	81.0	31.0
IV	S Cba	72.0	52.0	79.0	42.8
V	NCnt SFe	14.0	5.0	15.0	16.8
VI	North Belt	69.0	55.0	70.0	43.2
VII	South Belt	68.0	53.0	71.0	47.6
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	79.0	65.0	81.0	62.6
X	Cnt BA	77.0	40.0	75.0	59.4
XI	SW BA-S LP	-	-	-	-
XII	SE BA	67.0	60.0	70.0	55.4
XIII	SL	81.0	45.0	74.0	57.6
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		67.4	42.5	71.8	42.8

Corn

Corn Planted: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	55.0	21.0	30.0	36.8
II	NEA	38.0	23.0	37.0	37.6
III	NCnt Cba	96.0	93.9	100.0	93.6
IV	S Cba	98.0	97.0	100.0	96.0
V	NCnt SFe	61.0	67.0	75.8	73.2
VI	North Belt	99.0	100.0	100.0	99.4
VII	South Belt	99.0	100.0	100.0	99.4
VIII	ECnt ER	84.0	86.4	93.0	93.2
IX	N LP-W BA	98.0	97.2	100.0	98.8
X	Cnt BA	99.0	97.5	100.0	99.4
XI	SW BA-S LP	98.0	99.3	100.0	99.6
XII	SE BA	96.0	99.3	100.0	99.2
XIII	SL	98.0	94.1	100.0	95.6
XIV	Cuenca Sal	97.0	96.6	100.0	99.0
XV	Others	100.0	30.0	35.0	86.2
National		86.0	83.5	88.2	85.4

CROP CONDITION & DEVELOPMENT STAGES

Corn Condition: as of January 8, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	3.3	55.0	41.7	-
II	NEA	-	-	62.5	37.5	-
III	NCnt Cba	-	3.1	52.8	38.0	6.1
IV	S Cba	-	3.1	34.4	52.7	9.8
V	NCnt SFe	-	0.7	11.2	73.7	14.5
VI	North Belt	-	1.9	21.6	67.4	9.1
VII	South Belt	-	7.3	46.5	33.4	12.8
VIII	ECnt ER	-	-	60.0	38.0	2.0
IX	N LP-W BA	-	4.5	16.9	68.7	9.9
X	Cnt BA	-	5.8	22.4	68.8	3.0
XI	SW BA-S LP	5.0	33.2	38.3	22.1	1.4
XII	SE BA	-	7.2	71.8	21.1	-
XIII	SL	-	14.3	63.1	22.6	-
XIV	Cuenca Sal	1.8	16.8	28.3	47.3	6.0
XV	Others	-	10.0	50.0	40.0	-
National		0.2	5.3	39.7	48.0	6.8
Nat. Jan 9, 2019		0.4	5.5	43.0	43.0	8.1

Corn Moisture Condition: as of January 8, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	20.0	50.0	30.0	-
II	NEA	-	2.5	45.0	52.5	-
III	NCnt Cba	-	3.5	50.8	45.7	-
IV	S Cba	-	5.9	28.3	65.9	-
V	NCnt SFe	-	10.6	66.6	21.2	1.6
VI	North Belt	-	15.3	42.5	41.2	0.9
VII	South Belt	-	7.6	38.8	52.4	1.2
VIII	ECnt ER	-	0.6	31.9	67.2	0.3
IX	N LP-W BA	-	2.9	22.2	74.9	-
X	Cnt BA	0.4	7.4	40.0	52.2	-
XI	SW BA-S LP	15.0	57.0	23.1	4.9	-
XII	SE BA	5.8	29.6	36.2	28.5	-
XIII	SL	0.7	33.3	53.4	12.6	-
XIV	Cuenca Sal	7.2	17.8	25.5	49.5	-
XV	Others	-	-	50.0	50.0	-
National		0.9	10.9	38.8	49.1	0.3
Nat. Jan 9, 2019		0.0	9.2	49.7	34.8	6.3

CROP CONDITION & DEVELOPMENT STAGES

Corn Leaf Development: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	33.0	9.0	14.0	17.6
II	NEA	24.0	10.0	15.0	16.4
III	NCnt Cba	74.0	49.3	85.6	42.2
IV	S Cba	78.0	64.5	84.5	59.4
V	NCnt SFe	44.0	47.8	53.3	46.8
VI	North Belt	96.0	97.3	98.2	90.0
VII	South Belt	95.0	96.7	98.0	89.6
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	90.0	87.8	93.4	82.8
X	Cnt BA	87.0	78.4	91.0	78.0
XI	SW BA-S LP	-	-	-	-
XII	SE BA	84.0	86.0	89.5	88.4
XIII	SL	85.0	59.3	80.8	67.2
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		73.8	63.9	75.7	61.2

Sunflower

Sunflower Condition: as of January 8, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	-	-	-
II	NEA	-	-	38.0	58.0	4.0
III	NCnt Cba	-	-	-	-	-
IV	S Cba	-	-	-	-	-
V	NCnt SFe	-	-	32.0	63.0	5.0
VI	North Belt	-	-	-	-	-
VII	South Belt	-	-	-	-	-
VIII	ECnt ER	-	-	-	-	-
IX	N LP-W BA	-	3.0	40.0	52.0	5.0
X	Cnt BA	-	-	-	-	-
XI	SW BA-S LP	4.0	27.0	59.0	10.0	-
XII	SE BA	-	5.0	48.0	47.0	-
XIII	SL	-	-	-	-	-
XIV	Cuenca Sal	-	5.0	48.0	41.0	6.0
XV	Others	-	-	-	-	-
National		0.9	8.4	45.9	42.5	2.3
Nat. Jan 9, 2019		2.0	7.9	37.3	42.6	10.2

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Moisture Condition: as of January 8, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	-	-	-	-
II	NEA	-	-	37.0	59.0	4.0
III	NCnt Cba	-	-	-	-	-
IV	S Cba	-	-	-	-	-
V	NCnt SFe	-	-	35.0	60.0	5.0
VI	North Belt	-	-	-	-	-
VII	South Belt	-	-	-	-	-
VIII	ECnt ER	-	-	-	-	-
IX	N LP-W BA	-	2.0	45.0	53.0	-
X	Cnt BA	-	-	-	-	-
XI	SW BA-S LP	9.0	39.0	48.0	4.0	-
XII	SE BA	-	13.0	54.0	33.0	-
XIII	SL	-	-	-	-	-
XIV	Cuenca Sal	3.0	25.0	41.0	31.0	-
XV	Others	-	-	-	-	-
National		2.3	14.6	46.0	35.8	1.3
Nat. Jan 9, 2019		0.6	5.6	47.4	35.9	10.5

Sunflower Flowerbud Stage: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100.0	100.0	100.0	100.0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	100.0	100.0	100.0	100.0
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	86.0	79.0	92.0	93.8
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	50.0	40.0	63.0	62.4
XII	SE BA	51.0	42.0	58.0	66.2
XIII	SL	-	-	-	-
XIV	Cuenca Sal	83.0	74.0	88.0	85.6
XV	Others	-	-	-	-
National		76.2	68.7	80.1	81.5

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Flowering Stage: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100.0	100.0	100.0	100.0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	100.0	100.0	100.0	100.0
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	46.0	38.0	61.0	53.4
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	16.0	8.0	30.0	25.2
XII	SE BA	22.0	7.0	32.0	36.6
XIII	SL	-	-	-	-
XIV	Cuenca Sal	62.0	36.0	45.0	42.2
XV	Others	-	-	-	-
National		57.9	46.1	60.5	57.8

Sunflower Seed Development: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100.0	100.0	100.0	100.0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	100.0	96.0	100.0	97.6
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	15.0	-	7.0	15.4
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	1.8
XII	SE BA	-	-	-	2.6
XIII	SL	-	-	-	-
XIV	Cuenca Sal	13.0	-	-	10.4
XV	Others	-	-	-	-
National		41.9	34.1	36.2	36.5

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Mature: as of January 8, 2020

Regions	January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I NWA	-	-	-	-
II NEA	100.0	89.0	100.0	99.0
III NCnt Cba	-	-	-	-
IV S Cba	-	-	-	-
V NCnt SFe	76.0	62.0	85.0	76.4
VI North Belt	-	-	-	-
VII South Belt	-	-	-	-
VIII ECnt ER	-	-	-	-
IX N LP-W BA	-	-	-	-
X Cnt BA	-	-	-	-
XI SW BA-S LP	-	-	-	-
XII SE BA	-	-	-	-
XIII SL	-	-	-	-
XIV Cuenca Sal	-	-	-	-
XV Others	-	-	-	-
National	34.7	25.9	31.6	29.4

Wheat

Wheat Condition: as of January 8, 2020

Regions	Very Poor %	Poor %	Fair %	Good %	Excellent %
I NWA	-	-	-	-	-
II NEA	-	-	-	-	-
III NCnt Cba	-	-	-	-	-
IV S Cba	-	-	-	-	-
V NCnt SFe	-	-	-	-	-
VI North Belt	-	-	-	-	-
VII South Belt	-	-	-	-	-
VIII ECnt ER	-	-	-	-	-
IX N LP-W BA	-	-	-	-	-
X Cnt BA	-	-	-	-	-
XI SW BA-S LP	-	-	-	-	-
XII SE BA	4.0	46.0	50.0	-	-
XIII SL	-	-	-	-	-
XIV Cuenca Sal	-	-	-	-	-
XV Others	-	-	-	-	-
National	4.0	46.0	50.0	-	-
Nat. Jan 9, 2019	4.3	9.5	32.6	41.8	11.7

CROP CONDITION & DEVELOPMENT STAGES

Wheat Moisture Condition: as of January 8, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	-	-	-	-
II	NEA	-	-	-	-	-
III	NCnt Cba	-	-	-	-	-
IV	S Cba	-	-	-	-	-
V	NCnt SFe	-	-	-	-	-
VI	North Belt	-	-	-	-	-
VII	South Belt	-	-	-	-	-
VIII	ECnt ER	-	-	-	-	-
IX	N LP-W BA	-	-	-	-	-
X	Cnt BA	-	-	-	-	-
XI	SW BA-S LP	-	-	-	-	-
XII	SE BA	3.0	28.0	55.0	14.0	-
XIII	SL	-	-	-	-	-
XIV	Cuenca Sal	-	-	-	-	-
XV	Others	-	-	-	-	-
National		3.0	28.0	55.0	14.0	-
Nat. Jan 9, 2019		-	11.5	63.9	23.3	1.2

Wheat Mature: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	100.0	100.0	100.0	100.0
II	NEA	100.0	100.0	100.0	100.0
III	NCnt Cba	100.0	100.0	100.0	100.0
IV	S Cba	100.0	100.0	100.0	100.0
V	NCnt SFe	100.0	100.0	100.0	100.0
VI	North Belt	100.0	100.0	100.0	100.0
VII	South Belt	100.0	100.0	100.0	100.0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100.0	100.0	100.0	100.0
X	Cnt BA	100.0	100.0	100.0	99.6
XI	SW BA-S LP	100.0	95.0	100.0	99.6
XII	SE BA	100.0	82.0	100.0	98.0
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		100.0	97.8	100.0	99.7

CROP CONDITION & DEVELOPMENT STAGES

Wheat Harvested: as of January 8, 2020

Regions		January 9, 2019 %	January 1, 2020 %	January 8, 2020 %	2014/2019 Average %
I	NWA	100.0	100.0	100.0	100.0
II	NEA	100.0	100.0	100.0	100.0
III	NCnt Cba	100.0	100.0	100.0	100.0
IV	S Cba	100.0	100.0	100.0	100.0
V	NCnt SFe	100.0	100.0	100.0	100.0
VI	North Belt	100.0	100.0	100.0	100.0
VII	South Belt	100.0	100.0	100.0	100.0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100.0	100.0	100.0	99.0
X	Cnt BA	96.0	95.0	100.0	95.0
XI	SW BA-S LP	85.0	91.8	100.0	88.6
XII	SE BA	70.0	31.7	54.2	73.6
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		95.6	92.1	95.5	95.1