

CROP CONDITION & DEVELOPMENT STAGES


JANUARY, 2
2020



BUENOS AIRES GRAIN EXCHANGE
DEPARTMENT OF AGRICULTURAL
ESTIMATES

ISSN 2591-4715

DEPARTMENT & REGIONS



- I - NOA (Noroeste Argentino)
- II - NEA (Noreste Argentino)
- III - Centro-Norte de Córdoba
- IV - Sur de Córdoba
- V - Centro-Norte de Santa Fe
- VI - Núcleo Norte
- VII - Núcleo Sur
- VIII - Centro-Este de Entre Ríos
- IX - Norte de La Pampa - Oeste de Buenos Aires
- X - Centro de Buenos Aires
- XI - Sudoeste de Buenos Aires - Sur de La Pampa
- XII - Sudeste de Buenos Aires
- XIII - San Luis
- XIV - Cuenca del Salado
- XV - Corrientes - Misiones

HEAD OF DEPARTMENT

Esteban J Copati
ecopati@bc.org.ar

CROP ANALYST

Martin López
martinlopez@bc.org.ar

CROP ANALYST

Daniela A. Venturino
dventurino@bc.org.ar

CROP CONDITION ANALYST

Sofía Console Insúa
sconsoleinsua@bc.org.ar
Soybean

Matías Mihura
mmihura@bc.org.ar
Corn

Juan F. De Aramburu
jdearamburu@bc.org.ar
Wheat and Sunflower

CONTACT

123 Corrientes Avenue
C1043AAB - CABA
Phone: +54 11 4515 8200 | 8300
estimacionesagricolas@bc.org.ar
Twitter: @estimacionesbc

ISSN 2591-4715

CROP CONDITION & DEVELOPMENT STAGES

Findings of the survey conducted through January, 1 2020. The areas under analysis in this report account for 90% of the crops planted area. The regional development data is published once planting is complete. The national planting and harvest progress, as well as the phenological data of the crops derive from the final area projection, while the moisture and crop condition derive from the planting progress to date.

First Soybean

Season 2019/2020

First Soybean Planted: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	87,0	30,0	32,5	59,4
II	NEA	50,0	33,3	47,0	52,0
III	NCnt Cba	100,0	100,0	100,0	99,8
IV	S Cba	100,0	100,0	100,0	99,8
V	NCnt SFe	100,0	100,0	100,0	97,6
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	100,0	100,0	100,0	100,0
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	100,0	100,0	100,0	100,0
XI	SW BA-S LP	100,0	100,0	100,0	100,0
XII	SE BA	100,0	100,0	100,0	98,6
XIII	SL	100,0	95,4	100,0	100,0
XIV	Cuenca Sal	100,0	100,0	100,0	100,0
XV	Others	100,0	90,0	100,0	99,0
National		96,0	89,8	91,4	94,4

First Soybean Condition: as of January 1, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	14,0	53,0	33,0
II	NEA	-	-	17,0	52,0	31,0
III	NCnt Cba	-	3,0	48,0	49,0	-
IV	S Cba	-	3,0	55,0	42,0	-
V	NCnt SFe	-	-	24,0	64,0	12,0
VI	North Belt	-	-	51,0	43,0	6,0
VII	South Belt	-	-	25,0	71,0	4,0
VIII	ECnt ER	-	1,0	22,0	74,0	3,0
IX	N LP-W BA	-	2,0	70,0	28,0	-
X	Cnt BA	-	14,0	68,0	18,0	-
XI	SW BA-S LP	-	8,0	87,0	5,0	-
XII	SE BA	-	-	15,0	73,0	12,0
XIII	SL	-	-	80,0	20,0	-
XIV	Cuenca Sal	-	11,0	74,0	15,0	-
XV	Others	-	-	20,0	73,0	7,0
National		-	1,7	43,1	50,0	5,2
Nat. Jan 2, 2019		0,1	11,0	44,7	38,6	5,6

CROP CONDITION & DEVELOPMENT STAGES

First Soybean Moisture Condition: as of January 1, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	-	17,0	83,0	-
II	NEA	-	-	14,0	82,0	4,0
III	NCnt Cba	-	7,0	62,0	31,0	-
IV	S Cba	-	15,0	73,0	12,0	-
V	NCnt SFe	-	-	26,0	59,0	15,0
VI	North Belt	-	-	45,0	52,0	3,0
VII	South Belt	-	4,0	70,0	26,0	-
VIII	ECnt ER	-	-	40,0	57,0	3,0
IX	N LP-W BA	-	7,0	74,0	19,0	-
X	Cnt BA	1,0	22,0	60,0	17,0	-
XI	SW BA-S LP	3,0	35,0	58,0	4,0	-
XII	SE BA	-	16,0	56,0	28,0	-
XIII	SL	-	22,0	57,0	21,0	-
XIV	Cuenca Sal	1,0	31,0	66,0	2,0	-
XV	Others	-	-	27,0	63,0	10,0
National		0,1	6,5	55,5	35,9	2,0
Nat. Jan 2, 2019		-	6,3	54,7	32,6	6,4

First Soybean Leaf Development: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	73,0	74,0	90,0	82,6
IV	S Cba	85,0	90,0	98,0	88,4
V	NCnt SFe	73,0	96,0	100,0	74,6
VI	North Belt	95,0	100,0	100,0	87,6
VII	South Belt	96,0	100,0	100,0	95,4
VIII	ECnt ER	86,0	97,0	100,0	93,0
IX	N LP-W BA	90,0	97,0	100,0	92,8
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	71,0	90,0	100,0	76,8
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		78,5	80,2	86,3	80,0

CROP CONDITION & DEVELOPMENT STAGES

First Soybean Beginning Bloom: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	35,0	25,0	36,0	51,4
IV	S Cba	39,0	26,0	39,0	42,0
V	NCnt SFe	25,0	32,0	40,0	46,6
VI	North Belt	51,0	39,0	50,0	51,4
VII	South Belt	43,0	32,0	50,0	53,0
VIII	ECnt ER	27,0	23,0	37,0	48,6
IX	N LP-W BA	38,0	25,0	38,0	36,8
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	6,0	-	-	4,6
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		30,0	22,7	32,3	38,4

First Soybean Beginning Pod: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	5,0	-	-	11,2
IV	S Cba	-	-	-	6,2
V	NCnt SFe	-	-	1,0	7,6
VI	North Belt	8,0	-	-	7,8
VII	South Belt	-	-	-	3,6
VIII	ECnt ER	-	-	-	3,6
IX	N LP-W BA	-	-	-	2,0
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		1,6	-	0,1	4,2

CROP CONDITION & DEVELOPMENT STAGES

Second Soybean

Second Soybean Planted: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	45,0	7,0	10,0	48,0
II	NEA	35,0	15,0	40,7	45,0
III	NCnt Cba	97,0	77,0	90,3	94,4
IV	S Cba	98,0	65,0	87,6	92,6
V	NCnt SFe	94,0	68,0	86,0	85,8
VI	North Belt	98,0	100,0	100,0	97,2
VII	South Belt	85,0	80,0	90,0	91,0
VIII	ECnt ER	70,0	77,0	87,2	89,8
IX	N LP-W BA	80,0	70,8	82,4	87,8
X	Cnt BA	61,0	38,0	60,0	76,0
XI	SW BA-S LP	80,0	30,0	45,0	73,8
XII	SE BA	36,0	7,0	18,3	56,6
XIII	SL	75,0	60,0	86,6	85,6
XIV	Cuenca Sal	45,0	37,7	50,0	63,2
XV	Others	65,0	40,0	86,0	83,0
National		75,0	24,1	27,6	78,0

Second Soybean Condition: as of January 1, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	97,0	3,0	-
II	NEA	-	-	95,0	5,0	-
III	NCnt Cba	-	-	8,0	81,0	11,0
IV	S Cba	-	-	7,0	81,0	12,0
V	NCnt SFe	-	-	-	61,0	39,0
VI	North Belt	-	-	-	65,0	35,0
VII	South Belt	-	-	-	75,0	25,0
VIII	ECnt ER	-	-	-	57,0	43,0
IX	N LP-W BA	-	-	7,0	93,0	-
X	Cnt BA	-	-	91,0	9,0	-
XI	SW BA-S LP	-	-	13,0	87,0	-
XII	SE BA	-	-	55,0	45,0	-
XIII	SL	-	-	60,0	40,0	-
XIV	Cuenca Sal	-	-	50,0	50,0	-
XV	Others	-	-	42,0	58,0	-
National		-	-	14,8	66,0	19,2
Nat. Jan 2, 2019		0,3	12,0	55,4	31,8	0,5

CROP CONDITION & DEVELOPMENT STAGES

Second Soybean Moisture Condition: as of January 1, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	18,0	62,0	20,0	-
II	NEA	-	-	46,0	54,0	-
III	NCnt Cba	-	4,0	66,0	30,0	-
IV	S Cba	-	33,0	67,0	-	-
V	NCnt SFe	-	8,0	92,0	-	-
VI	North Belt	-	17,0	56,0	27,0	-
VII	South Belt	-	7,0	78,0	15,0	-
VIII	ECnt ER	-	7,0	85,0	8,0	-
IX	N LP-W BA	-	30,0	70,0	-	-
X	Cnt BA	-	39,0	61,0	-	-
XI	SW BA-S LP	-	8,0	52,0	40,0	-
XII	SE BA	-	5,0	53,0	42,0	-
XIII	SL	12,0	48,0	40,0	-	-
XIV	Cuenca Sal	-	-	52,0	48,0	-
XV	Others	-	-	46,0	54,0	-
National		0,0	15,2	67,4	17,3	-
Nat.Jan 2, 2019		-	8,2	51,1	33,0	7,8

Second Soybean Leaf Development: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	-	-	-	-
VI	North Belt	59,0	-	60,0	61,6
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		46,4	-	31,7	35,7

CROP CONDITION & DEVELOPMENT STAGES

Soybean

Soybean Planted: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	20,0	2,0	7,0	24,4
II	NEA	25,0	-	15,0	36,0
III	NCnt Cba	85,0	45,1	77,0	77,8
IV	S Cba	85,0	37,4	65,0	80,2
V	NCnt SFe	75,0	60,5	68,0	72,0
VI	North Belt	93,0	80,3	100,0	90,2
VII	South Belt	72,0	64,0	80,0	81,4
VIII	ECnt ER	65,0	70,3	77,0	74,6
IX	N LP-W BA	60,0	29,6	70,8	71,6
X	Cnt BA	30,0	30,0	38,0	49,4
XI	SW BA-S LP	40,0	9,0	30,0	39,8
XII	SE BA	30,0	-	7,0	31,6
XIII	SL	55,0	-	60,0	73,6
XIV	Cuenca Sal	15,0	-	37,7	42,0
XV	Others	40,0	-	40,0	65,0
National		61,0	18,2	24,1	60,6

Soybean Condition: as of January 1, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	100,0	-	-
II	NEA	-	-	100,0	-	-
III	NCnt Cba	-	-	10,0	80,0	10,0
IV	S Cba	-	-	10,0	80,0	10,0
V	NCnt SFe	-	-	-	60,0	40,0
VI	North Belt	-	-	-	70,0	30,0
VII	South Belt	-	-	-	80,0	20,0
VIII	ECnt ER	-	-	-	60,0	40,0
IX	N LP-W BA	-	-	10,0	90,0	-
X	Cnt BA	-	-	100,0	-	-
XI	SW BA-S LP	-	-	15,0	85,0	-
XII	SE BA	-	-	60,0	40,0	-
XIII	SL	-	-	55,0	45,0	-
XIV	Cuenca Sal	-	-	49,0	51,0	-
XV	Others	-	-	45,0	55,0	-
National		-	-	11,9	69,3	18,7
Nat. Jan 2, 2019		1,3	15,2	60,6	22,7	0,3

CROP CONDITION & DEVELOPMENT STAGES

Soybean Moisture Condition: as of January 1, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	20,0	60,0	20,0	-
II	NEA	-	-	50,0	50,0	-
III	NCnt Cba	-	8,0	63,0	29,0	-
IV	S Cba	-	36,0	64,0	-	-
V	NCnt SFe	-	10,0	90,0	-	-
VI	North Belt	-	30,0	50,0	20,0	-
VII	South Belt	-	11,0	76,0	13,0	-
VIII	ECnt ER	-	10,0	84,0	6,0	-
IX	N LP-W BA	-	40,0	60,0	-	-
X	Cnt BA	-	50,0	50,0	-	-
XI	SW BA-S LP	10,0	50,0	40,0	-	-
XII	SE BA	-	10,0	50,0	40,0	-
XIII	SL	10,0	45,0	45,0	-	-
XIV	Cuenca Sal	-	15,0	60,0	25,0	-
XV	Others	-	-	50,0	50,0	-
National		0,3	23,3	63,5	12,9	-
Nat. Jan 2, 2019		-	9,2	57,4	26,8	6,6

Soybean Leaf Development: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	-	-	-	-
VI	North Belt	83,7	65,1	65,1	82,0
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		-	-	-	-

CROP CONDITION & DEVELOPMENT STAGES

Early Corn

Early Corn Planted: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	42,0	5,0	21,0	28,2
II	NEA	33,0	4,0	23,0	28,6
III	NCnt Cba	100,0	100,0	100,0	100,0
IV	S Cba	100,0	100,0	100,0	100,0
V	NCnt SFe	100,0	100,0	100,0	100,0
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	100,0	100,0	100,0	100,0
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	100,0	100,0	100,0	100,0
XI	SW BA-S LP	100,0	100,0	100,0	100,0
XII	SE BA	100,0	100,0	100,0	100,0
XIII	SL	100,0	100,0	100,0	100,0
XIV	Cuenca Sal	100,0	100,0	100,0	100,0
XV	Others	100,0	26,0	30,0	100,0
National		79,0	75,5	80,0	85,6

Early Corn Condition: as of January 1, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	55,0	45,0	-
II	NEA	-	-	43,0	57,0	-
III	NCnt Cba	-	8,0	62,0	29,0	1,0
IV	S Cba	-	12,0	53,0	35,0	-
V	NCnt SFe	-	-	36,0	56,0	8,0
VI	North Belt	-	18,0	46,0	36,0	-
VII	South Belt	-	10,0	43,0	45,0	2,0
VIII	ECnt ER	-	8,0	42,0	42,0	8,0
IX	N LP-W BA	-	14,0	64,0	22,0	-
X	Cnt BA	3,0	29,0	53,0	15,0	-
XI	SW BA-S LP	-	17,0	64,0	19,0	-
XII	SE BA	-	17,0	50,0	33,0	-
XIII	SL	-	19,0	46,0	35,0	-
XIV	Cuenca Sal	-	28,0	53,0	19,0	-
XV	Others	-	15,0	60,0	25,0	-
National		0,2	13,5	50,5	34,7	1,2
Nat. Jan 2, 2019		0,3	5,1	38,6	47,5	8,5

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Moisture Condition: as of January 1, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	4,0	36,0	60,0	-
II	NEA	-	1,0	36,0	63,0	-
III	NCnt Cba	-	35,0	47,0	18,0	-
IV	S Cba	-	29,0	55,0	16,0	-
V	NCnt SFe	-	1,0	26,0	66,0	7,0
VI	North Belt	-	35,0	48,0	17,0	-
VII	South Belt	-	35,0	43,0	22,0	-
VIII	ECnt ER	-	12,0	37,0	46,0	5,0
IX	N LP-W BA	1,0	23,0	56,0	18,0	2,0
X	Cnt BA	2,0	59,0	30,0	9,0	-
XI	SW BA-S LP	4,0	43,0	38,0	15,0	-
XII	SE BA	2,0	50,0	36,0	12,0	-
XIII	SL	-	31,0	55,0	14,0	-
XIV	Cuenca Sal	2,0	47,0	44,0	7,0	-
XV	Others	-	-	45,0	55,0	-
National		0,5	30,7	44,4	23,6	0,8
Nat. Jan 2, 2019		0,2	9,1	50,6	34,9	5,2

Early Corn Leaf Development: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	10,0	-	9,0	10,2
II	NEA	9,0	-	10,0	8,4
III	NCnt Cba	100,0	100,0	100,0	85,0
IV	S Cba	100,0	100,0	100,0	92,0
V	NCnt SFe	100,0	100,0	100,0	91,4
VI	North Belt	100,0	100,0	100,0	96,8
VII	South Belt	100,0	100,0	100,0	96,8
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100,0	100,0	100,0	97,4
X	Cnt BA	100,0	100,0	100,0	99,0
XI	SW BA-S LP	-	-	-	-
XII	SE BA	100,0	100,0	100,0	98,0
XIII	SL	100,0	100,0	100,0	95,2
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		74,1	74,4	76,8	69,8

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Pollination: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	100,0	70,0	85,0	76,2
IV	S Cba	100,0	69,0	82,0	84,4
V	NCnt SFe	100,0	100,0	100,0	91,4
VI	North Belt	100,0	95,0	99,0	96,2
VII	South Belt	100,0	85,0	96,0	93,4
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	85,0	64,0	82,0	80,2
X	Cnt BA	71,0	55,0	72,0	73,0
XI	SW BA-S LP	-	-	-	-
XII	SE BA	16,0	14,0	20,0	19,0
XIII	SL	70,0	25,0	40,0	52,2
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		63,2	54,1	61,7	55,8

Early Corn Silking: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	84,0	52,0	70,0	64,0
IV	S Cba	82,0	51,0	60,0	72,8
V	NCnt SFe	100,0	96,0	99,0	91,4
VI	North Belt	98,0	85,0	98,0	88,2
VII	South Belt	93,0	69,0	86,0	84,2
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	63,0	45,0	64,0	66,6
X	Cnt BA	49,0	22,0	49,0	50,0
XI	SW BA-S LP	-	-	-	-
XII	SE BA	6,0	3,0	14,0	5,0
XIII	SL	35,0	10,0	29,0	26,2
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		55,8	42,8	53,8	48,0

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Grain Growth: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	5,0	-	4,0	3,6
IV	S Cba	1,0	-	1,0	1,2
V	NCnt SFe	59,0	35,0	51,0	48,6
VI	North Belt	14,0	7,0	14,0	15,0
VII	South Belt	9,0	-	7,0	6,0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		8,2	2,8	6,4	5,8

Late Corn

Late Corn Planted: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	90,0	81,0	92,0	81,2
IV	S Cba	95,0	83,0	96,0	86,4
V	NCnt SFe	35,0	6,0	40,0	36,6
VI	North Belt	90,0	68,0	100,0	86,4
VII	South Belt	90,0	77,0	100,0	85,6
VIII	ECnt ER	38,0	30,0	61,0	62,6
IX	N LP-W BA	92,0	80,0	92,0	89,8
X	Cnt BA	95,0	81,0	93,0	95,6
XI	SW BA-S LP	90,0	80,0	98,0	92,4
XII	SE BA	88,0	82,0	98,0	90,8
XIII	SL	95,0	80,0	92,0	87,8
XIV	Cuenca Sal	90,0	75,0	91,0	93,4
XV	Others	-	-	-	-
National		87,0	74,4	89,3	74,2

CROP CONDITION & DEVELOPMENT STAGES

Late Corn Condition: as of January 1, 2020

Regions	Very Poor %	Poor %	Fair %	Good %	Excellent %
I NWA	-	-	-	-	-
II NEA	-	-	-	-	-
III NCnt Cba	-	5,0	60,0	34,0	1,0
IV S Cba	-	3,0	58,0	33,0	6,0
V NCnt SFe	-	-	25,0	63,0	12,0
VI North Belt	-	4,0	51,0	38,0	7,0
VII South Belt	-	4,0	49,0	43,0	4,0
VIII ECnt ER	-	-	35,0	60,0	5,0
IX N LP-W BA	-	25,0	63,0	10,0	2,0
X Cnt BA	-	24,0	58,0	18,0	-
XI SW BA-S LP	-	18,0	60,0	20,0	2,0
XII SE BA	-	8,0	34,0	58,0	-
XIII SL	-	20,0	60,0	20,0	-
XIV Cuenca Sal	-	35,0	53,0	12,0	-
XV Others	-	-	-	-	-
National	-	8,9	55,7	32,0	3,4
Nat.Jan 2, 2019	1,0	6,3	54,0	35,5	3,2

Late Corn Moisture Condition: as of January 1, 2020

Regions	Dry %	Poor %	Favourable %	Optimum %	Wet %
I NWA	-	-	-	-	-
II NEA	-	-	-	-	-
III NCnt Cba	-	19,0	63,0	18,0	-
IV S Cba	-	12,0	53,0	35,0	-
V NCnt SFe	-	-	35,0	60,0	5,0
VI North Belt	-	40,0	44,0	16,0	-
VII South Belt	-	39,0	43,0	18,0	-
VIII ECnt ER	-	-	32,0	58,0	10,0
IX N LP-W BA	1,0	41,0	41,0	17,0	-
X Cnt BA	4,0	50,0	36,0	10,0	-
XI SW BA-S LP	5,0	46,0	41,0	8,0	-
XII SE BA	1,0	55,0	39,0	5,0	-
XIII SL	-	30,0	50,0	20,0	-
XIV Cuenca Sal	1,0	59,0	35,0	5,0	-
XV Others	-	-	-	-	-
National	0,4	26,9	49,9	22,4	0,4
Nat.Jan 2, 2019	2,5	27,3	49,0	17,3	3,9

CROP CONDITION & DEVELOPMENT STAGES

Late Corn Leaf Development: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	-	-	-	-
VI	North Belt	34,0	14,0	55,0	20,2
VII	South Belt	51,0	18,0	53,0	26,4
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		39,0	10,8	42,5	21,2

Corn

Corn Planted: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	42,0	5,0	21,0	25,2
II	NEA	33,0	4,0	23,0	24,8
III	NCnt Cba	92,0	85,6	93,9	84,0
IV	S Cba	96,0	87,4	97,0	90,4
V	NCnt SFe	58,0	48,3	67,0	59,6
VI	North Belt	99,0	98,1	100,0	97,4
VII	South Belt	98,0	98,4	100,0	96,6
VIII	ECnt ER	75,0	75,5	86,4	86,4
IX	N LP-W BA	96,0	93,0	97,2	95,0
X	Cnt BA	97,0	93,2	97,5	97,8
XI	SW BA-S LP	95,0	92,6	99,3	97,2
XII	SE BA	94,0	93,7	99,3	97,2
XIII	SL	96,0	85,2	94,1	90,8
XIV	Cuenca Sal	94,0	91,3	96,6	97,0
XV	Others	100,0	26,0	30,0	80,8
National		83,0	75,1	83,5	79,0

CROP CONDITION & DEVELOPMENT STAGES

Corn Condition: as of January 1, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	55,0	45,0	-
II	NEA	-	-	43,0	57,0	-
III	NCnt Cba	-	5,8	60,5	32,7	1,0
IV	S Cba	-	5,4	56,7	33,5	4,4
V	NCnt SFe	-	-	32,4	58,3	9,3
VI	North Belt	-	12,6	47,9	36,8	2,7
VII	South Belt	-	9,6	43,4	44,9	2,1
VIII	ECnt ER	-	6,0	40,3	46,5	7,3
IX	N LP-W BA	-	17,6	63,7	18,0	0,7
X	Cnt BA	2,0	27,3	54,7	16,0	-
XI	SW BA-S LP	-	17,4	62,5	19,4	0,7
XII	SE BA	-	13,9	44,5	41,6	-
XIII	SL	-	19,7	56,1	24,1	-
XIV	Cuenca Sal	-	30,3	53,0	16,7	-
XV	Others	-	15,0	60,0	25,0	-
National		0,1	11,5	52,8	33,5	2,2
Nat.Jan 2, 2019		0,6	5,7	45,9	41,8	6,0

Corn Moisture Condition: as of January 1, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	4,0	36,0	60,0	-
II	NEA	-	1,0	36,0	63,0	-
III	NCnt Cba	-	23,1	58,9	18,0	-
IV	S Cba	-	16,6	53,5	29,9	-
V	NCnt SFe	-	0,7	29,0	64,0	6,3
VI	North Belt	-	36,9	46,4	16,6	-
VII	South Belt	-	35,3	43,0	21,7	-
VIII	ECnt ER	-	9,0	35,8	49,0	6,2
IX	N LP-W BA	1,0	29,0	51,0	17,7	1,3
X	Cnt BA	2,7	55,9	32,1	9,3	-
XI	SW BA-S LP	4,4	44,1	39,1	12,4	-
XII	SE BA	1,7	51,7	37,0	9,6	-
XIII	SL	-	30,3	51,4	18,3	-
XIV	Cuenca Sal	1,7	50,9	41,0	6,3	-
XV	Others	-	-	45,0	55,0	-
National		0,5	29,0	46,8	23,1	0,6
Nat.Jan 2, 2019		1,3	17,8	49,8	26,5	4,6

CROP CONDITION & DEVELOPMENT STAGES

Corn Leaf Development: as of January 1, 2020

Regions	January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I NWA	-	-	-	-
II NEA	-	-	-	-
III NCnt Cba	-	-	-	-
IV S Cba	-	-	-	-
V NCnt SFe	-	-	-	-
VI North Belt	91,0	94,8	97,3	86,0
VII South Belt	93,0	94,3	96,7	85,8
VIII ECnt ER	-	-	-	-
IX N LP-W BA	-	-	-	-
X Cnt BA	-	-	-	-
XI SW BA-S LP	-	-	-	-
XII SE BA	-	-	-	-
XIII SL	-	-	-	-
XIV Cuenca Sal	-	-	-	-
XV Others	-	-	-	-
National	57,9	50,5	63,9	50,8

Sunflower

Sunflower Condition: as of January 1, 2020

Regions	Very Poor %	Poor %	Fair %	Good %	Excellent %
I NWA	-	-	-	-	-
II NEA	-	-	34,0	56,0	10,0
III NCnt Cba	-	-	-	-	-
IV S Cba	-	-	-	-	-
V NCnt SFe	-	7,0	34,0	51,0	8,0
VI North Belt	-	-	-	-	-
VII South Belt	-	-	-	-	-
VIII ECnt ER	-	-	-	-	-
IX N LP-W BA	-	5,0	54,0	41,0	-
X Cnt BA	-	-	-	-	-
XI SW BA-S LP	-	21,0	67,0	12,0	-
XII SE BA	-	-	44,0	54,0	2,0
XIII SL	-	-	-	-	-
XIV Cuenca Sal	-	10,0	54,0	36,0	-
XV Others	-	-	-	-	-
National	-	6,9	47,7	41,8	3,5
Nat. Jan 2, 2019	3,3	7,8	40,8	40,3	7,8

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Moisture Condition: as of January 1, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	-	-	-	-
II	NEA	-	-	39,0	55,0	6,0
III	NCnt Cba	-	-	-	-	-
IV	S Cba	-	-	-	-	-
V	NCnt SFe	-	-	34,0	57,0	9,0
VI	North Belt	-	-	-	-	-
VII	South Belt	-	-	-	-	-
VIII	ECnt ER	-	-	-	-	-
IX	N LP-W BA	-	7,0	53,0	40,0	-
X	Cnt BA	-	-	-	-	-
XI	SW BA-S LP	3,0	25,0	63,0	9,0	-
XII	SE BA	-	-	56,0	44,0	-
XIII	SL	-	-	-	-	-
XIV	Cuenca Sal	5,0	23,0	52,0	20,0	-
XV	Others	-	-	-	-	-
National		1,0	7,9	50,9	37,8	2,4
Nat. Jan 2, 2019		0,9	8,6	54,2	28,9	7,4

Sunflower Flowerbud Stage: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100,0	100,0	100,0	100,0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	100,0	100,0	100,0	100,0
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	71,0	60,0	79,0	85,2
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	31,0	21,0	40,0	46,4
XII	SE BA	33,0	23,0	42,0	49,8
XIII	SL	-	-	-	-
XIV	Cuenca Sal	72,0	60,0	74,0	75,4
XV	Others	-	-	-	-
National		66,4	57,1	68,7	72,2

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Flowering Stage: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100,0	100,0	100,0	100,0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	100,0	100,0	100,0	100,0
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	29,0	25,0	38,0	39,4
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	8,0	10,8
XII	SE BA	4,0	-	7,0	18,6
XIII	SL	-	-	-	-
XIV	Cuenca Sal	48,0	18,0	36,0	27,6
XV	Others	-	-	-	-
National		47,8	39,6	46,1	47,4

Sunflower Seed Development: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100,0	100,0	100,0	100,0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	100,0	84,0	96,0	95,0
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		39,2	31,9	34,1	33,0

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Mature: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	92,0	74,0	89,0	91,6
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	63,0	49,0	62,0	62,6
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		30,7	21,2	25,9	26,0

Wheat

Wheat Condition: as of January 1, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	-	-	-
II	NEA	-	-	-	-	-
III	NCnt Cba	-	-	-	-	-
IV	S Cba	-	-	-	-	-
V	NCnt SFe	-	-	-	-	-
VI	North Belt	-	-	-	-	-
VII	South Belt	-	-	-	-	-
VIII	ECnt ER	-	-	-	-	-
IX	N LP-W BA	-	-	-	-	-
X	Cnt BA	27,0	64,0	9,0	-	-
XI	SW BA-S LP	56,0	44,0	-	-	-
XII	SE BA	11,0	50,0	39,0	-	-
XIII	SL	-	-	-	-	-
XIV	Cuenca Sal	-	-	-	-	-
XV	Others	-	-	-	-	-
National		15,4	49,8	34,8	-	-
Nat. Jan 2, 2019		4,6	13,8	38,3	37,5	5,8

CROP CONDITION & DEVELOPMENT STAGES

Wheat Moisture Condition: as of January 1, 2020

Regions	Dry %	Poor %	Favourable %	Optimum %	Wet %
I NWA	-	-	-	-	-
II NEA	-	-	-	-	-
III NCnt Cba	-	-	-	-	-
IV S Cba	-	-	-	-	-
V NCnt SFe	-	-	-	-	-
VI North Belt	-	-	-	-	-
VII South Belt	-	-	-	-	-
VIII ECnt ER	-	-	-	-	-
IX N LP-W BA	-	-	-	-	-
X Cnt BA	14,0	37,0	49,0	-	-
XI SW BA-S LP	41,0	53,0	6,0	-	-
XII SE BA	8,0	31,0	57,0	4,0	-
XIII SL	-	-	-	-	-
XIV Cuenca Sal	-	-	-	-	-
XV Others	-	-	-	-	-
National	10,9	33,3	52,1	3,7	-
Nat.Jan 2, 2019	-	9,4	53,1	27,0	10,5

Wheat Mature: as of January 1, 2020

Regions	January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I NWA	100,0	100,0	100,0	100,0
II NEA	100,0	100,0	100,0	100,0
III NCnt Cba	100,0	100,0	100,0	100,0
IV S Cba	100,0	100,0	100,0	100,0
V NCnt SFe	100,0	100,0	100,0	100,0
VI North Belt	100,0	100,0	100,0	100,0
VII South Belt	100,0	100,0	100,0	100,0
VIII ECnt ER	-	-	-	-
IX N LP-W BA	100,0	96,0	100,0	100,0
X Cnt BA	100,0	95,0	100,0	95,4
XI SW BA-S LP	84,0	75,0	95,0	89,8
XII SE BA	83,0	64,0	82,0	84,2
XIII SL	-	-	-	-
XIV Cuenca Sal	-	-	-	-
XV Others	-	-	-	-
National	97,0	93,5	97,8	96,8

CROP CONDITION & DEVELOPMENT STAGES

Wheat Harvested: as of January 1, 2020

Regions		January 2, 2019 %	December 25, 2019 %	January 1, 2020 %	2014/2019 Average %
I	NWA	100,0	100,0	100,0	100,0
II	NEA	100,0	100,0	100,0	100,0
III	NCnt Cba	100,0	100,0	100,0	100,0
IV	S Cba	100,0	100,0	100,0	100,0
V	NCnt SFe	100,0	100,0	100,0	100,0
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	96,9	100,0	100,0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	85,0	94,0	100,0	91,4
X	Cnt BA	78,0	91,7	95,0	81,0
XI	SW BA-S LP	70,0	73,3	91,8	69,4
XII	SE BA	55,0	22,5	31,7	48,2
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		90,6	87,8	92,1	88,2