



DEPARTAMENT & REGIONS

CROP ANALYST

Ing. Martin López martinlopez @bc.org.ar

CROP ANALYST

Ing. Daniela A. Venturino dventurino @bc.org.ar

CROP ANALYST

Ing. Andrés Paterniti apaterniti @bc.org.ar

CROP SURVEY & ANALYSIS OF CROP CONDITION AND DEVELOPMENT STAGES

Joaquín Pellejero jpellejero @bc.org.ar


Agustín Podestá apodesta @bc.org.ar

CONTACT

Av. Corrientes 123

C1043AAB - CABA
Tel.: +54 11 3221 7230
estimacionesagricolas@bc.org.ar
Twitter: @BolsadeC_ETyM
www.bolsadecereales.com

ISSN 2408-4344t



I - NWA (North-West Argentina)

II - NEA (North-East Argentina)

III - North-Central Córdoba

IV - South Córdoba

V - North-Central Santa Fe

VI - North Belt

VII - South Belt

VIII - East-Central Entre Ríos

IX - North La Pampa - West Buenos

X - Central Buenos Aires

XI - South-West de Buenos Aires - South La Pampa

XII - South-East Buenos Aires

XIII - San Luis


XIV - Cuenca del Salado

XV - Others

We appreciate the contribution of our Network of Collaborators throughout the country.



RECENT ADVERSE EVENTS





METHODOLOGY

Findings of the survey conducted through October 12, 2022. The areas under analysis in this report account for 90% of the crops planted area. The regional development data is published once planting is complete. The national planting and harvest progress, as well as the phenological data of the crops derive from the final area projection, while the moisture and crop condition derive from the planting progress to



SUNFLOWER 2022/23












Regional participation in the total national area (%)															
Regions	XII	ΧI	V	IX	II.	XIV	Х	IV	XIII	VI	VII	VIII	XV	III	1
Nat. Part. 2022/23	26,3	23,7	13,2	9,0	8,5	6,1	4,7	2,6	2,4	1,3	0,6	0,5	0,4	0,9	0,0
Hist. Nat. Part.	25,1	22,6	13,2	7,8	15,0	6,3	4,1	1,6	1,5	0,8	0,5	0,5	0,5	0,4	0,0
Moisture o	Moisture condition, Crop condition & Development stages Week 42														
Regions	Adverse event		Season		Moisture Condition Favourable / Optimum		Crop Condition Fair / Excellent		Planted	Leaves Development	Flowerbud Stage	Flowering stage	Seed development	Mature	Harvested
			22/23		55,0		62,0		100,0	95,0	32,0	7,0	-	-	-
II - NEA			21/22 17/22 Average		35,0 -		100,0		100,0 100,0	100,0 100,0	43,0 40,7	6,0 5,3	-	-	-
V - NC S Fe	Frost		22/23		54,0		85,0		100,0	65,0	14,0	-	-	-	-
			21/22		38,0		94,0		100,0	95,0	19,0	-	-	-	-
			17/22 Average		-		-		100,0	95,2	19,8	-	-	-	-
IX - N LP -			22/23		92,0				7,0	-	-	-	-	-	-
W BA			21/22		75,0		-		27,5	-	-	-	-	-	-
			17/22 Average		-		-		13,2	-	-	-	-	-	-
X - Central BA		22/23		/23	90,0		-		5,0	-	-	-	-	-	-
			21/22		100,0			-		-	-	-	-	-	-
				22 Average		-		-	12,0	0,5	-	-	-	-	-
XI - SW BA -			22/23		100,0		-		6,0	-	-	-	-	-	-
SLP			21/22		94,0		-		11,2	-	-	-	-	-	-
			17/22 Average		-		-		5,9	-	-	-	-	-	-
XII - SE BA			22/23		96,0		-		5,0	-	-	-	-	-	-
			21/22		97,0		-		11,3	-	-	-	-	-	-
			17/22 Average		-		-		8,7	-	-	-	-	-	-
XIV - Cuenca Sal.			22/23		80,0		-		5,0	-	-	-	-	-	-
			21/22		100,0		-		12,5	-	-	-	-	-	-
				Average		-		-	19,0	-	-	-		-	-
National				/23		58,1		79,5	29,6	18,1	4,6	0,6	-	-	-
				/22		54,8		97,0	38,9	26,0	7,0	0,6	-	-	-
			17/22 Average		-		-		39,1	30,7	9,3	1,0	-	-	-



WHEAT 2022/23









Regional participation in the total national area (%)															
Regions	XII	VI	VII	ΧI	IX	V	III.	III	VIII	X	IV	1	XIV	XV	XIII
Nat. Part. 2022/23	13,5	12,6	12,1	11,0	9,6	7,2	6,9	5,4	6,8	6,4	4,4	3,0	0,8	0,4	0,2
Hist. Nat. Part.	10,4	13,3	10,0	9,2	8,3	7,6	6,9	9,3	5,1	5,5	6,2	6,2	1,6	0,3	0,2
Moisture o	isture condition, Crop condition & Development stages Week 42														42
Regions	Adverse event		Season		Moisture Condition Favourable / Optimum		Crop Condition Fair / Excellent		Planted	Tillering	Steam Extension	Heading	Ripening	Mature	Harvested
	Drought		22/23			10,0	11,0		100,0	100,0	100,0	100,0	68,0	40,0	-
I - NWA			21/22		-		-		100,0	100,0	100,0	100,0	80,0	48,0	-
			17/22 Average						100,0	100,0	100,0	100,0	71,8	39,0	-
	Drought		22/23 23		23,0	14,0		100,0	100,0	100,0	100,0	42,0	21,0	-	
II - NEA			21,	21/22 15,0		5,0		100,0	100,0	100,0	100,0	68,0	41,0	-	
			17/22 Average		-	-		100,0	100,0	100,0	100,0	50,6	22,8	-	
III - NC Cba	Drought		22/	/23	21,0		12,0		100,0	100,0	100,0	62,0	5,0	-	-
			21/22		35,0		39,0	100,0	100,0	100,0	73,0	9,0	-	-	
			17/22 Average		-		-		100,0	100,0	99,4	68,6	8,4	-	-
IV - S Cba	Drought		22/23		30,0		23,0		100,0	100,0	100,0	77,0	-	-	
			21/22		55,0		62,0		100,0	100,0	100,0	50,0	-	-	-
			17/22	Average	-		-		100,0	100,0	97,8	48,2	0,6	-	-
V - NC S Fe	Drought		22/	/23	40,0		34,0		100,0	100,0	100,0	72,0	15,0	1,0	
			21/	/22	47,0		57,0		100,0	100,0	100,0	76,0	20,0	4,0	1,2
			17/22	Average	-		-		100,0	100,0	100,0	75,4	19,8	1,4	0,2
VI - North Belt	Drought - Frost		22/	/23	24,0		18,0		100,0	100,0	94,0	49,0	-	-	
			21/	/22	85,0		100,0		100,0	100,0	97,0	60,0	7,0	-	-
			17/22	Average	-		-		100,0	100,0	99,2	61,2	4,6	0,2	-
VII - South Belt	Drought - Frost		22/	/23	17,0		22,0		100,0	100,0	84,0	41,0	-	-	
			21/	/22	22 85,0			100,0	100,0	100,0	93,0	47,0	1,0	-	-
			17/22	Average	-		-		100,0	100,0	93,6	44,4	0,2	-	-
IX - N LP - W BA			22/	/23		62,0		70,0	100,0	100,0	58,0	17,0	-	-	-
			21,	/22	74,0		100,0		100,0	100,0	79,0	32,0	-	-	-
			17/22	Average		-		-	100,0	100,0	75,0	24,8	_	_	_
X - Central BA			22/	/23		61,0		87,0	100,0	100,0	57,0	11,0	-	-	-
			21,	/22	96,0		100,0		100,0	100,0	76,0	7,0	-	-	-
			17/22	Average	I				100,0	100,0	66,6	17,2	-	-	-
VI CILLER			22/	/23		74,0		88,0	100,0	100,0	40,0	9,0	-	-	-
S LP				/22		71,0		89,0	100,0	100,0	60,0	10,0	-	-	-
			17/22	Average		-		-	100,0	99,4	56,6	11,6			
XII - SE BA	Frost		22/	/23	85,0		91,0		100,0	100,0	41,0	3,0	-	-	-
			21,	/22	95,0		100,0		100,0	100,0	57,0	13,0	-	-	-
			ı	Average	1 1		-		100,0	100,0	56,2	13,8	_	_	-
National				/23		47,0		50,9	100,0	100,0	75,5	40,7	6,4	2,7	1,7
				/22		68,1		77,9	100,0	100,0	85,9	48,4	12,2	5,7	-
			ı	Average		-		-	100,0	100,0	85,8	49,3	11,0	4,1	-