

CROP CONDITION & DEVELOPMENT STAGES


NOVEMBER, 1
2018



BUENOS AIRES GRAIN EXCHANGE
DEPARTMENT OF AGRICULTURAL
ESTIMATES

ISSN 2591-4715

DEPARTMENT & REGIONS



- I - NOA (Noroeste Argentino)
- II - NEA (Noreste Argentino)
- III - Centro-Norte de Córdoba
- IV - Sur de Córdoba
- V - Centro-Norte de Santa Fe
- VI - Núcleo Norte
- VII - Núcleo Sur
- VIII - Centro-Este de Entre Ríos
- IX - Norte de La Pampa - Oeste de Buenos Aires
- X - Centro de Buenos Aires
- XI - Sudoeste de Buenos Aires - Sur de La Pampa
- XII - Sudeste de Buenos Aires
- XIII - San Luis
- XIV - Cuenca del Salado
- XV - Corrientes - Misiones

HEAD OF DEPARTMENT

Esteban J Copati
ecopati@bc.org.ar

CROP ANALYST

Martin López
martinlopez@bc.org.ar

COORDINATOR

Gonzalo Hermida
ghermida@bc.org.ar

SATELLITE IMAGE ANALYST

María Victoria Corte
mcorte@bc.org.ar

CROP CONDITION ANALYST

Daniela A. Venturino
dventurino@bc.org.ar
Wheat & Sunflower

Santiago Burrone
sburrone@bc.org.ar
Corn

Ayelén Gago
amgago@bc.org.ar
Soybean

CONTACT

123 Corrientes Avenue
C1043AAB - CABA
Phone: +54 11 4515 8200 | 8300
estimacionesagricolas@bc.org.ar
Twitter: @estimacionesbc

ISSN 2591-4715

CROP CONDITION & DEVELOPMENT STAGES

Findings of the survey conducted through October 31, 2018. The areas under analysis in this report account for 90% of the crops planted area. The regional development data is published once planting is complete. The national planting and harvest progress, as well as the phenological data of the crops derive from the final area projection, while the moisture and crop condition derive from the planting progress to date.

Early Corn

Early Corn Planted: as of October 31, 2018

| Regions | | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|--------------|------------------------|-----------------------|-----------------------|------------------------|
| I | NOA | - | - | - | - |
| II | NEA | - | - | - | - |
| III | Ctro N Cba | 100,0 | 97,0 | 100,0 | 90,0 |
| IV | S Cba | 100,0 | 98,0 | 100,0 | 88,0 |
| V | Ctro N SFe | 100,0 | 100,0 | 100,0 | 96,0 |
| VI | Núcleo Norte | 100,0 | 100,0 | 100,0 | 98,0 |
| VII | Núcleo Sur | 100,0 | 100,0 | 100,0 | 99,0 |
| VIII | Ctro E ER | 100,0 | 100,0 | 100,0 | 100,0 |
| IX | N LP-OBA | 95,0 | 95,0 | 100,0 | 97,0 |
| X | Ctro Bs As | 90,0 | 84,0 | 88,0 | 91,8 |
| XI | SO BA-S LP | 75,0 | 70,0 | 85,0 | 65,4 |
| XII | SE BA | 50,0 | 42,0 | 60,0 | 51,2 |
| XIII | San Luis | 100,0 | 100,0 | 100,0 | 72,0 |
| XIV | Cuenca Sal | 95,0 | 90,0 | 95,0 | 82,2 |
| XV | Others | 100,0 | 100,0 | 100,0 | 100,0 |
| National | | 62,8 | 62,7 | 64,7 | 62,6 |

Early Corn Condition: as of October 31, 2018

| Regions | | Very Poor % | Poor % | Fair % | Good % | Excellent % |
|---------------------------|--------------|----------------|-------------|-------------|-------------|----------------|
| I | NOA | - | - | - | - | - |
| II | NEA | - | - | - | - | - |
| III | Ctro N Cba | 0,5 | 16,5 | 49,5 | 28,7 | 4,8 |
| IV | S Cba | - | 13,7 | 59,1 | 24,1 | 3,1 |
| V | Ctro N SFe | - | 7,0 | 34,1 | 55,4 | 3,5 |
| VI | Núcleo Norte | - | 12,4 | 60,2 | 24,9 | 2,5 |
| VII | Núcleo Sur | - | 12,5 | 40,1 | 41,3 | 6,1 |
| VIII | Ctro E ER | - | - | - | - | - |
| IX | N LP-OBA | - | 21,2 | 51,9 | 22,2 | 4,7 |
| X | Ctro Bs As | 24,1 | 28,7 | 33,1 | 12,1 | 2,0 |
| XI | SO BA-S LP | - | - | - | - | - |
| XII | SE BA | 1,1 | 22,1 | 46,9 | 29,1 | 0,8 |
| XIII | San Luis | 2,1 | 41,5 | 40,8 | 15,6 | - |
| XIV | Cuenca Sal | - | - | - | - | - |
| XV | Others | - | - | - | - | - |
| National | | 1,8 | 16,2 | 48,5 | 29,5 | 4,0 |
| Nat. Nov. 01, 2017 | | - | 1,9 | 16,4 | 76,3 | 5,4 |

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Moisture Condition: as of October 31, 2018

| Regions | Dry % | Poor % | Favourable % | Optimum % | Wet % |
|--------------------------|------------|-------------|--------------|-------------|-------------|
| I NOA | - | - | - | - | - |
| II NEA | - | - | - | - | - |
| III Ctro N Cba | 2,2 | 54,5 | 43,3 | - | - |
| IV S Cba | 1,2 | 37,8 | 45,7 | 14,2 | 1,1 |
| V Ctro N SFe | - | 28,2 | 55,1 | 16,7 | - |
| VI Núcleo Norte | - | 35,2 | 58,0 | 6,8 | - |
| VII Núcleo Sur | - | 16,4 | 54,3 | 17,9 | 11,4 |
| VIII Ctro E ER | - | - | - | - | - |
| IX N LP-OBA | - | 6,8 | 58,7 | 27,5 | 7,0 |
| X Ctro Bs As | - | 9,0 | 60,1 | 16,0 | 14,9 |
| XI SO BA-S LP | - | - | - | - | - |
| XII SE BA | 1,1 | 17,8 | 55,0 | 25,5 | 0,6 |
| XIII San Luis | 10,1 | 57,9 | 28,6 | 3,4 | - |
| XIV Cuenca Sal | - | - | - | - | - |
| XV Others | - | - | - | - | - |
| National | 0,9 | 26,6 | 53,8 | 13,9 | 4,8 |
| Nat.Nov. 01, 2017 | - | 4,6 | 26,6 | 40,4 | 28,5 |

Early Corn Leaf Development: as of October 31, 2018

| Regions | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|---------------------|--------------------|--------------------|---------------------|
| I NOA | - | - | - | - |
| II NEA | - | - | - | - |
| III Ctro N Cba | 88,9 | 70,1 | 84,4 | 64,0 |
| IV S Cba | 88,9 | 75,3 | 88,2 | 66,9 |
| V Ctro N SFe | 100,0 | 100,0 | 100,0 | 92,0 |
| VI Núcleo Norte | 100,0 | 87,4 | 94,0 | 86,8 |
| VII Núcleo Sur | 95,0 | 82,0 | 92,5 | 84,2 |
| VIII Ctro E ER | - | - | - | - |
| IX N LP-OBA | 70,0 | 52,4 | 74,9 | 68,5 |
| X Ctro Bs As | - | - | - | - |
| XI SO BA-S LP | - | - | - | - |
| XII SE BA | - | - | - | - |
| XIII San Luis | 91,0 | 43,1 | 68,1 | 58,2 |
| XIV Cuenca Sal | - | - | - | - |
| XV Others | - | - | - | - |
| National | 54,9 | 43,6 | 51,8 | 49,0 |

CROP CONDITION & DEVELOPMENT STAGES

Corn

Corn Planted: as of October 31, 2018

| Regions | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|------------------------|-----------------------|-----------------------|------------------------|
| I NOA | - | - | - | - |
| II NEA | - | - | - | - |
| III Ctro N Cba | 15,0 | 14,6 | 15,0 | 12,3 |
| IV S Cba | 20,0 | 19,6 | 20,0 | 24,8 |
| V Ctro N SFe | 35,0 | 35,0 | 35,0 | 34,8 |
| VI Núcleo Norte | 85,0 | 87,0 | 87,0 | 76,6 |
| VII Núcleo Sur | 80,0 | 85,0 | 85,0 | 75,3 |
| VIII Ctro E ER | 60,0 | 60,0 | 60,0 | 62,0 |
| IX N LP-OBA | 47,5 | 47,5 | 50,0 | 51,4 |
| X Ctro Bs As | 36,0 | 37,8 | 39,6 | 49,1 |
| XI SO BA-S LP | 45,0 | 35,0 | 42,5 | 45,5 |
| XII SE BA | 30,0 | 21,0 | 30,0 | 40,3 |
| XIII San Luis | 20,0 | 20,0 | 20,0 | 14,7 |
| XIV Cuenca Sal | 38,0 | 40,5 | 42,8 | 49,9 |
| XV Others | 40,0 | 100,0 | 100,0 | 38,0 |
| National | 33,9 | 34,4 | 35,5 | 35,8 |

Sunflower

Sunflower Planted: as of October 31, 2018

| Regions | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|------------------------|-----------------------|-----------------------|------------------------|
| I NOA | - | - | - | - |
| II NEA | 100,0 | 100,0 | 100,0 | 100,0 |
| III Ctro N Cba | 100,0 | 100,0 | 100,0 | 71,0 |
| IV S Cba | 30,0 | 70,0 | 90,0 | 43,4 |
| V Ctro N SFe | 100,0 | 100,0 | 100,0 | 100,0 |
| VI Núcleo Norte | 92,0 | 100,0 | 100,0 | 82,0 |
| VII Núcleo Sur | 25,0 | 50,0 | 75,0 | 57,0 |
| VIII Ctro E ER | 85,0 | 100,0 | 100,0 | 91,0 |
| IX N LP-OBA | 14,0 | 26,0 | 45,0 | 50,0 |
| X Ctro Bs As | 12,0 | 30,0 | 42,0 | 56,2 |
| XI SO BA-S LP | 8,0 | 22,0 | 30,0 | 33,0 |
| XII SE BA | 10,0 | 25,0 | 40,0 | 34,6 |
| XIII San Luis | 15,0 | 20,0 | 50,0 | 51,0 |
| XIV Cuenca Sal | 25,0 | 44,0 | 60,0 | 57,0 |
| XV Others | 100,0 | 100,0 | 100,0 | 85,0 |
| National | 44,6 | 56,0 | 64,2 | 58,5 |

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Condition: as of October 31, 2018

| Regions | | Very Poor % | Poor % | Fair % | Good % | Excellent % |
|--------------------------|--------------|-------------|------------|-------------|-------------|-------------|
| I | NOA | - | - | - | - | - |
| II | NEA | - | - | 26,0 | 52,0 | 22,0 |
| III | Ctro N Cba | - | - | - | - | - |
| IV | S Cba | - | - | - | - | - |
| V | Ctro N SFe | - | 8,0 | 33,0 | 52,0 | 7,0 |
| VI | Núcleo Norte | - | - | - | - | - |
| VII | Núcleo Sur | - | - | - | - | - |
| VIII | Ctro E ER | - | - | - | - | - |
| IX | N LP-OBA | - | 4,0 | 42,0 | 38,0 | 16,0 |
| X | Ctro Bs As | - | - | - | - | - |
| XI | SO BA-S LP | - | - | 71,0 | 29,0 | - |
| XII | SE BA | - | 6,0 | 76,0 | 18,0 | - |
| XIII | San Luis | - | - | - | - | - |
| XIV | Cuenca Sal | - | 11,0 | 36,0 | 48,0 | 5,0 |
| XV | Others | - | - | - | - | - |
| National | | 0,1 | 4,6 | 41,1 | 43,4 | 10,9 |
| Nat.Nov. 01, 2017 | | - | 3,4 | 12,4 | 80,6 | 3,7 |

Sunflower Moisture Condition: as of October 31, 2018

| Regions | | Dry % | Poor % | Favourable % | Optimum % | Wet % |
|--------------------------|--------------|------------|-------------|--------------|-------------|-------------|
| I | NOA | - | - | - | - | - |
| II | NEA | - | 9,0 | 61,0 | 27,0 | 3,0 |
| III | Ctro N Cba | - | - | - | - | - |
| IV | S Cba | - | - | - | - | - |
| V | Ctro N SFe | - | 39,0 | 48,0 | 13,0 | - |
| VI | Núcleo Norte | - | - | - | - | - |
| VII | Núcleo Sur | - | - | - | - | - |
| VIII | Ctro E ER | - | - | - | - | - |
| IX | N LP-OBA | - | 12,0 | 45,0 | 35,0 | 8,0 |
| X | Ctro Bs As | - | - | - | - | - |
| XI | SO BA-S LP | - | 12,0 | 51,0 | 31,0 | 6,0 |
| XII | SE BA | 3,0 | 35,0 | 41,0 | 21,0 | - |
| XIII | San Luis | - | - | - | - | - |
| XIV | Cuenca Sal | - | 15,0 | 55,0 | 30,0 | - |
| XV | Others | - | - | - | - | - |
| National | | 1,0 | 21,5 | 51,6 | 23,3 | 2,6 |
| Nat.Nov. 01, 2017 | | - | 12,4 | 40,6 | 30,9 | 16,0 |

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Leaf Development: as of October 31, 2018

| Regions | | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|--------------|------------------------|-----------------------|-----------------------|------------------------|
| I | NOA | - | - | - | - |
| II | NEA | 100,0 | 100,0 | 100,0 | 100,0 |
| III | Ctro N Cba | - | - | - | - |
| IV | S Cba | - | - | - | - |
| V | Ctro N SFe | 100,0 | 100,0 | 100,0 | 100,0 |
| VI | Núcleo Norte | - | - | - | - |
| VII | Núcleo Sur | - | - | - | - |
| VIII | Ctro E ER | - | - | - | - |
| IX | N LP-OBA | 18,0 | 12,0 | 22,0 | 25,0 |
| X | Ctro Bs As | - | - | - | - |
| XI | SO BA-S LP | 14,0 | 2,0 | 12,0 | 8,8 |
| XII | SE BA | 15,0 | 2,0 | 15,0 | 10,2 |
| XIII | San Luis | - | - | - | - |
| XIV | Cuenca Sal | 19,0 | 6,0 | 20,0 | 20,8 |
| XV | Others | - | - | - | - |
| National | | 48,0 | 42,1 | 49,4 | 39,8 |

Sunflower Flowerbud Stage: as of October 31, 2018

| Regions | | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|--------------|------------------------|-----------------------|-----------------------|------------------------|
| I | NOA | - | - | - | - |
| II | NEA | 75,0 | 61,0 | 75,0 | 68,0 |
| III | Ctro N Cba | - | - | - | - |
| IV | S Cba | - | - | - | - |
| V | Ctro N SFe | 46,0 | 32,0 | 41,0 | 45,6 |
| VI | Núcleo Norte | - | - | - | - |
| VII | Núcleo Sur | - | - | - | - |
| VIII | Ctro E ER | - | - | - | - |
| IX | N LP-OBA | - | - | - | - |
| X | Ctro Bs As | - | - | - | - |
| XI | SO BA-S LP | - | - | - | - |
| XII | SE BA | - | - | - | - |
| XIII | San Luis | - | - | - | - |
| XIV | Cuenca Sal | - | - | - | - |
| XV | Others | - | - | - | - |
| National | | 23,3 | 19,2 | 24,1 | 17,9 |

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Flowering Stage: as of October 31, 2018

| Regions | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|------------------------|-----------------------|-----------------------|------------------------|
| I NOA | - | - | - | - |
| II NEA | 29,0 | 27,0 | 42,0 | 36,4 |
| III Ctro N Cba | - | - | - | - |
| IV S Cba | - | - | - | - |
| V Ctro N SFe | 5,0 | - | 9,0 | 10,6 |
| VI Núcleo Norte | - | - | - | - |
| VII Núcleo Sur | - | - | - | - |
| VIII Ctro E ER | - | - | - | - |
| IX N LP-OBA | - | - | - | - |
| X Ctro Bs As | - | - | - | - |
| XI SO BA-S LP | - | - | - | - |
| XII SE BA | - | - | - | - |
| XIII San Luis | - | - | - | - |
| XIV Cuenca Sal | - | - | - | - |
| XV Others | - | - | - | - |
| National | 7,0 | 6,3 | 11,2 | 7,8 |

Sunflower Seed Development: as of October 31, 2018

| Regions | November 01, 2017 % | October 24, 2018 % | October 31, 2018 % | 2013/2018 Average % |
|-----------------|------------------------|-----------------------|-----------------------|------------------------|
| I NOA | - | - | - | - |
| II NEA | - | 15,0 | 23,0 | 9,2 |
| III Ctro N Cba | - | - | - | - |
| IV S Cba | - | - | - | - |
| V Ctro N SFe | - | - | - | - |
| VI Núcleo Norte | - | - | - | - |
| VII Núcleo Sur | - | - | - | - |
| VIII Ctro E ER | - | - | - | - |
| IX N LP-OBA | - | - | - | - |
| X Ctro Bs As | - | - | - | - |
| XI SO BA-S LP | - | - | - | - |
| XII SE BA | - | - | - | - |
| XIII San Luis | - | - | - | - |
| XIV Cuenca Sal | - | - | - | - |
| XV Others | - | - | - | - |
| National | - | 3,5 | 5,3 | 1,6 |

CROP CONDITION & DEVELOPMENT STAGES

Wheat

Wheat Condition: as of October 31, 2018

| Regions | | Very Poor % | Poor % | Fair % | Good % | Excellent % |
|--------------------------|--------------|----------------|-------------|-------------|-------------|----------------|
| I | NOA | 16,0 | 27,0 | 45,0 | 12,0 | - |
| II | NEA | - | 6,0 | 36,0 | 38,0 | 20,0 |
| III | Ctro N Cba | 3,0 | 34,0 | 38,0 | 17,0 | 8,0 |
| IV | S Cba | 4,0 | 46,0 | 37,0 | 9,0 | 4,0 |
| V | Ctro N SFe | 3,0 | 24,0 | 36,0 | 31,0 | 6,0 |
| VI | Núcleo Norte | 16,0 | 32,0 | 37,0 | 15,0 | - |
| VII | Núcleo Sur | 4,0 | 10,0 | 40,0 | 38,0 | 8,0 |
| VIII | Ctro E ER | - | - | - | - | - |
| IX | N LP-OBA | - | 6,0 | 33,0 | 54,0 | 7,0 |
| X | Ctro Bs As | - | 5,0 | 18,0 | 55,0 | 22,0 |
| XI | SO BA-S LP | - | 2,0 | 36,0 | 46,0 | 16,0 |
| XII | SE BA | - | 6,0 | 35,0 | 46,0 | 13,0 |
| XIII | San Luis | - | - | - | - | - |
| XIV | Cuenca Sal | - | - | - | - | - |
| XV | Others | - | - | - | - | - |
| National | | 4,4 | 18,0 | 36,3 | 32,5 | 8,7 |
| Nat.Nov. 01, 2017 | | 5,7 | 18,1 | 25,1 | 30,3 | 20,8 |

Wheat Moisture Condition: as of October 31, 2018

| Regions | | Dry % | Poor % | Favourable % | Optimum % | Wet % |
|--------------------------|--------------|------------|-------------|-----------------|--------------|-------------|
| I | NOA | - | 17,0 | 36,0 | 25,0 | 22,0 |
| II | NEA | - | 14,0 | 54,0 | 28,0 | 4,0 |
| III | Ctro N Cba | - | 58,0 | 33,0 | 9,0 | - |
| IV | S Cba | 25,0 | 46,0 | 21,0 | 8,0 | - |
| V | Ctro N SFe | 6,0 | 44,0 | 37,0 | 11,0 | 2,0 |
| VI | Núcleo Norte | 9,0 | 45,0 | 35,0 | 11,0 | - |
| VII | Núcleo Sur | - | 24,0 | 45,0 | 22,0 | 9,0 |
| VIII | Ctro E ER | - | - | - | - | - |
| IX | N LP-OBA | - | 6,0 | 61,0 | 28,0 | 5,0 |
| X | Ctro Bs As | - | 10,0 | 35,0 | 38,0 | 17,0 |
| XI | SO BA-S LP | 2,0 | 16,0 | 41,0 | 33,0 | 8,0 |
| XII | SE BA | 7,0 | 23,0 | 44,0 | 26,0 | - |
| XIII | San Luis | - | - | - | - | - |
| XIV | Cuenca Sal | - | - | - | - | - |
| XV | Others | - | - | - | - | - |
| National | | 4,9 | 30,2 | 39,8 | 20,5 | 4,6 |
| Nat.Nov. 01, 2017 | | 2,3 | 9,8 | 33,3 | 34,3 | 20,3 |

CROP CONDITION & DEVELOPMENT STAGES

Wheat Steam Extension: as of October 31, 2018

| Regions | | November 01, 2017 | October 24, 2018 | October 31, 2018 | 2013/2018 Average |
|-----------------|--------------|-------------------|------------------|------------------|-------------------|
| | | % | % | % | % |
| I | NOA | 100,0 | 100,0 | 100,0 | 100,0 |
| II | NEA | 100,0 | 100,0 | 100,0 | 100,0 |
| III | Ctro N Cba | 100,0 | 100,0 | 100,0 | 100,0 |
| IV | S Cba | 100,0 | 100,0 | 100,0 | 100,0 |
| V | Ctro N SFe | 100,0 | 100,0 | 100,0 | 100,0 |
| VI | Núcleo Norte | 100,0 | 100,0 | 100,0 | 100,0 |
| VII | Núcleo Sur | 100,0 | 100,0 | 100,0 | 100,0 |
| VIII | Ctro E ER | - | - | - | - |
| IX | N LP-OBA | 100,0 | 94,0 | 100,0 | 100,0 |
| X | Ctro Bs As | 85,0 | 91,0 | 97,0 | 97,0 |
| XI | SO BA-S LP | 70,0 | 81,0 | 95,0 | 89,4 |
| XII | SE BA | 69,0 | 92,0 | 100,0 | 86,6 |
| XIII | San Luis | - | - | - | - |
| XIV | Cuenca Sal | - | - | - | - |
| XV | Others | - | - | - | - |
| National | | 92,9 | 96,6 | 99,4 | 96,8 |

Wheat Heading: as of October 31, 2018

| Regions | | November 01, 2017 | October 24, 2018 | October 31, 2018 | 2013/2018 Average |
|-----------------|--------------|-------------------|------------------|------------------|-------------------|
| | | % | % | % | % |
| I | NOA | 100,0 | 100,0 | 100,0 | 100,0 |
| II | NEA | 100,0 | 100,0 | 100,0 | 100,0 |
| III | Ctro N Cba | 89,0 | 92,0 | 100,0 | 96,4 |
| IV | S Cba | 65,0 | 86,0 | 98,0 | 82,4 |
| V | Ctro N SFe | 100,0 | 85,0 | 96,0 | 100,0 |
| VI | Núcleo Norte | 75,0 | 100,0 | 100,0 | 89,8 |
| VII | Núcleo Sur | 64,0 | 72,0 | 85,0 | 85,8 |
| VIII | Ctro E ER | - | - | - | - |
| IX | N LP-OBA | 40,0 | 48,0 | 61,0 | 77,4 |
| X | Ctro Bs As | 41,0 | 52,0 | 65,0 | 65,0 |
| XI | SO BA-S LP | 39,0 | 39,0 | 54,0 | 62,6 |
| XII | SE BA | 29,0 | 44,0 | 57,0 | 60,2 |
| XIII | San Luis | - | - | - | - |
| XIV | Cuenca Sal | - | - | - | - |
| XV | Others | - | - | - | - |
| National | | 67,2 | 75,5 | 84,2 | 82,2 |

CROP CONDITION & DEVELOPMENT STAGES

Wheat Ripening: as of October 31, 2018

| Regions | | November 01, 2017 | October 24, 2018 | October 31, 2018 | 2013/2018 Average |
|-----------------|--------------|-------------------|------------------|------------------|-------------------|
| | | % | % | % | % |
| I | NOA | 90,0 | 88,0 | 100,0 | 87,2 |
| II | NEA | 70,0 | 69,0 | 84,0 | 78,6 |
| III | Ctro N Cba | 47,0 | 32,0 | 46,0 | 55,2 |
| IV | S Cba | 16,0 | 19,0 | 35,0 | 33,6 |
| V | Ctro N SFe | 68,0 | 35,0 | 48,0 | 68,2 |
| VI | Núcleo Norte | 41,0 | 23,0 | 35,0 | 52,4 |
| VII | Núcleo Sur | 15,0 | 6,0 | 19,0 | 27,8 |
| VIII | Ctro E ER | - | - | - | - |
| IX | N LP-OBA | - | - | - | 18,6 |
| X | Ctro Bs As | - | - | - | 11,2 |
| XI | SO BA-S LP | - | - | - | 16,0 |
| XII | SE BA | - | - | - | 12,0 |
| XIII | San Luis | - | - | - | - |
| XIV | Cuenca Sal | - | - | - | - |
| XV | Others | - | - | - | - |
| National | | 30,8 | 23,4 | 32,3 | 38,6 |

Wheat Mature: as of October 31, 2018

| Regions | | November 01, 2017 | October 24, 2018 | October 31, 2018 | 2013/2018 Average |
|-----------------|--------------|-------------------|------------------|------------------|-------------------|
| | | % | % | % | % |
| I | NOA | 66,0 | 61,0 | 71,0 | 60,0 |
| II | NEA | 42,0 | 43,0 | 57,0 | 55,0 |
| III | Ctro N Cba | 3,0 | - | 8,0 | 18,2 |
| IV | S Cba | - | 5,0 | 13,0 | 10,0 |
| V | Ctro N SFe | 19,0 | 10,0 | 22,0 | 32,0 |
| VI | Núcleo Norte | 7,0 | - | - | 15,6 |
| VII | Núcleo Sur | - | - | - | 4,4 |
| VIII | Ctro E ER | - | - | - | - |
| IX | N LP-OBA | - | - | - | 4,0 |
| X | Ctro Bs As | - | - | - | 3,6 |
| XI | SO BA-S LP | - | - | - | 2,8 |
| XII | SE BA | - | - | - | 1,8 |
| XIII | San Luis | - | - | - | - |
| XIV | Cuenca Sal | - | - | - | - |
| XV | Others | - | - | - | - |
| National | | 9,7 | 8,3 | 12,6 | 14,6 |