

CROP CONDITION & DEVELOPMENT STAGES


JANUARY, 16
2020



BUENOS AIRES GRAIN EXCHANGE
DEPARTMENT OF AGRICULTURAL
ESTIMATES

ISSN 2591-4715

DEPARTMENT & REGIONS



- I - NOA (Noroeste Argentino)
- II - NEA (Noreste Argentino)
- III - Centro-Norte de Córdoba
- IV - Sur de Córdoba
- V - Centro-Norte de Santa Fe
- VI - Núcleo Norte
- VII - Núcleo Sur
- VIII - Centro-Este de Entre Ríos
- IX - Norte de La Pampa - Oeste de Buenos Aires
- X - Centro de Buenos Aires
- XI - Sudoeste de Buenos Aires - Sur de La Pampa
- XII - Sudeste de Buenos Aires
- XIII - San Luis
- XIV - Cuenca del Salado
- XV - Corrientes - Misiones

HEAD OF DEPARTMENT

Esteban J Copati
ecopati@bc.org.ar

CROP ANALYST

Martin López
martinlopez@bc.org.ar

CROP ANALYST

Daniela A. Venturino
dventurino@bc.org.ar

CROP CONDITION ANALYST

Sofía Console Insúa
sconsoleinsua@bc.org.ar
Soybean, Wheat, Sunflower

Matías Mihura
mmihura@bc.org.ar
Corn

CONTACT

123 Corrientes Avenue
C1043AAB - CABA
Phone: +54 11 4515 8200 | 8300
estimacionesagricolas@bc.org.ar
Twitter: @estimacionesbc

ISSN 2591-4715

CROP CONDITION & DEVELOPMENT STAGES

Findings of the survey conducted through January, 8 2020. The areas under analysis in this report account for 90% of the crops planted area. The regional development data is published once planting is complete. The national planting and harvest progress, as well as the phenological data of the crops derive from the final area projection, while the moisture and crop condition derive from the planting progress to date.

First Soybean

Season 2019/2020

First Soybean Planted: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	96,0	50,0	58,6	86,0
II	NEA	86,0	78,9	82,0	85,0
III	NCnt Cba	100,0	100,0	100,0	100,0
IV	S Cba	100,0	100,0	100,0	100,0
V	NCnt SFe	100,0	100,0	100,0	100,0
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	100,0	100,0	100,0	100,0
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	100,0	100,0	100,0	100,0
XI	SW BA-S LP	100,0	100,0	100,0	100,0
XII	SE BA	100,0	100,0	100,0	100,0
XIII	SL	100,0	100,0	100,0	100,0
XIV	Cuenca Sal	100,0	100,0	100,0	100,0
XV	Others	100,0	100,0	100,0	100,0
National		99,0	95,3	96,1	98,0

First Soybean Condition: as of January 15, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	15,0	60,0	25,0
II	NEA	-	-	13,0	59,0	28,0
III	NCnt Cba	-	1,0	52,0	47,0	-
IV	S Cba	-	-	56,0	44,0	-
V	NCnt SFe	-	-	17,0	69,0	14,0
VI	North Belt	-	-	46,0	48,0	6,0
VII	South Belt	-	-	23,0	74,0	3,0
VIII	ECnt ER	-	9,0	69,0	22,0	-
IX	N LP-W BA	-	-	70,0	30,0	-
X	Cnt BA	-	9,0	70,0	21,0	-
XI	SW BA-S LP	-	10,0	87,0	3,0	-
XII	SE BA	-	-	23,0	69,0	8,0
XIII	SL	-	1,0	78,0	21,0	-
XIV	Cuenca Sal	-	6,0	76,0	18,0	-
XV	Others	-	-	18,0	73,0	9,0
National		-	1,3	44,7	48,5	5,6
Nat. Jan 16, 2019		1,6	7,1	37,1	41,6	12,5

CROP CONDITION & DEVELOPMENT STAGES

First Soybean Moisture Condition: as of January 15, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	-	33,0	67,0	-
II	NEA	-	-	14,0	86,0	-
III	NCnt Cba	-	2,0	65,0	33,0	-
IV	S Cba	-	7,0	77,0	15,0	1,0
V	NCnt SFe	-	-	26,0	66,0	8,0
VI	North Belt	-	-	41,0	56,0	3,0
VII	South Belt	-	1,0	63,0	35,0	1,0
VIII	ECnt ER	-	-	35,0	64,0	1,0
IX	N LP-W BA	-	2,0	68,0	30,0	-
X	Cnt BA	-	17,0	66,0	17,0	-
XI	SW BA-S LP	3,0	43,0	52,0	2,0	-
XII	SE BA	-	17,0	60,0	23,0	-
XIII	SL	2,0	21,0	62,0	15,0	-
XIV	Cuenca Sal	-	22,0	67,0	11,0	-
XV	Others	-	-	30,0	66,0	4,0
National		0,1	3,6	52,5	42,6	1,3
Nat. Jan 16, 2019		-	0,6	35,8	46,5	17,2

First Soybean Leaf Development: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	95,0	76,0	100,0	96,6
IV	S Cba	100,0	100,0	100,0	98,8
V	NCnt SFe	96,0	100,0	100,0	95,6
VI	North Belt	100,0	100,0	100,0	97,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	100,0	100,0	100,0	100,0
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	93,0	100,0	100,0	94,2
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		92,6	87,7	92,1	92,0

CROP CONDITION & DEVELOPMENT STAGES

First Soybean Beginning Bloom: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	56,0	48,0	56,0	72,6
IV	S Cba	60,0	58,0	71,0	68,4
V	NCnt SFe	50,0	50,0	60,0	71,0
VI	North Belt	72,0	68,0	83,0	76,2
VII	South Belt	65,0	70,0	84,0	78,8
VIII	ECnt ER	49,0	47,0	56,0	71,8
IX	N LP-W BA	62,0	56,0	76,0	64,0
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	25,0	10,0	32,0	27,2
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		50,4	46,4	58,6	60,9

First Soybean Beginning Pod: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	26,0	10,0	28,0	31,8
IV	S Cba	22,0	18,0	27,0	31,6
V	NCnt SFe	12,0	5,0	14,0	29,4
VI	North Belt	28,0	12,0	31,0	32,6
VII	South Belt	18,0	6,0	29,0	25,2
VIII	ECnt ER	5,0	-	-	17,0
IX	N LP-W BA	17,0	10,0	19,0	16,4
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		14,8	7,1	17,8	20,9

CROP CONDITION & DEVELOPMENT STAGES

First Soybean Beginning Seed: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	-	-	-	3,2
IV	S Cba	-	-	-	-
V	NCnt SFe	-	2,0	5,0	-
VI	North Belt	-	-	-	1,6
VII	South Belt	-	-	-	1,0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		-	0,1	0,3	0,8

Second Soybean

Second Soybean Planted: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	95,0	38,0	60,0	83,2
II	NEA	90,0	80,6	82,0	80,2
III	NCnt Cba	100,0	98,0	100,0	99,8
IV	S Cba	100,0	100,0	100,0	100,0
V	NCnt SFe	100,0	93,2	100,0	96,4
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	92,0	100,0	100,0	97,8
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	95,0	83,9	100,0	99,0
XI	SW BA-S LP	100,0	70,0	81,0	98,0
XII	SE BA	100,0	67,0	80,0	98,0
XIII	SL	100,0	100,0	100,0	100,0
XIV	Cuenca Sal	90,0	65,0	65,0	97,0
XV	Others	100,0	90,0	90,0	100,0
National		98,0	88,7	93,6	96,6

CROP CONDITION & DEVELOPMENT STAGES

Second Soybean Condition: as of January 15, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	96,0	4,0	-
II	NEA	-	-	88,0	12,0	-
III	NCnt Cba	-	-	6,0	81,0	13,0
IV	S Cba	-	-	1,0	85,0	14,0
V	NCnt SFe	-	-	-	61,0	39,0
VI	North Belt	-	-	-	63,0	37,0
VII	South Belt	-	-	2,0	75,0	23,0
VIII	ECnt ER	-	-	5,0	59,0	36,0
IX	N LP-W BA	-	-	3,0	92,0	5,0
X	Cnt BA	-	-	83,0	17,0	-
XI	SW BA-S LP	-	2,0	11,0	87,0	-
XII	SE BA	-	-	51,0	49,0	-
XIII	SL	-	-	60,0	40,0	-
XIV	Cuenca Sal	-	-	43,0	57,0	-
XV	Others	-	-	40,0	60,0	-
National		-	0,1	20,8	62,0	17,1
Nat. Jan 16, 2019		2,7	9,8	43,6	38,1	5,8

Second Soybean Moisture Condition: as of January 15, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	15,0	63,0	22,0	-
II	NEA	-	-	39,0	61,0	-
III	NCnt Cba	-	-	63,0	37,0	-
IV	S Cba	-	15,0	70,0	15,0	-
V	NCnt SFe	-	4,0	88,0	7,0	1,0
VI	North Belt	-	-	56,0	43,0	1,0
VII	South Belt	-	2,0	76,0	21,0	1,0
VIII	ECnt ER	-	-	84,0	13,0	3,0
IX	N LP-W BA	-	16,0	71,0	13,0	-
X	Cnt BA	-	24,0	69,0	7,0	-
XI	SW BA-S LP	-	13,0	58,0	29,0	-
XII	SE BA	-	-	58,0	42,0	-
XIII	SL	8,0	54,0	38,0	-	-
XIV	Cuenca Sal	-	-	43,0	57,0	-
XV	Others	-	-	48,0	52,0	-
National		0,0	6,4	66,2	26,8	0,5
Nat. Jan 16, 2019		-	0,9	36,1	45,7	17,4

CROP CONDITION & DEVELOPMENT STAGES

Second Soybean Leaf Development: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	82,0	56,0	84,0	73,6
IV	S Cba	73,0	66,0	76,0	69,4
V	NCnt SFe	75,0	50,0	77,0	69,4
VI	North Belt	90,0	73,0	91,0	87,8
VII	South Belt	76,0	50,0	77,0	68,6
VIII	ECnt ER	75,0	60,0	79,0	77,0
IX	N LP-W BA	80,0	50,0	84,0	72,8
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		70,6	48,1	72,7	61,2

Second Soybean Beginning Bloom: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	15,0	1,0	7,0	8,6
IV	S Cba	14,0	-	5,0	4,8
V	NCnt SFe	9,0	1,0	4,0	3,8
VI	North Belt	16,0	-	18,0	18,6
VII	South Belt	14,0	-	9,0	10,8
VIII	ECnt ER	-	-	-	3,8
IX	N LP-W BA	2,0	-	1,0	2,8
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		7,6	0,5	5,7	5,1

CROP CONDITION & DEVELOPMENT STAGES

Soybean

Soybean Planted: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	96,0	45,8	59,1	85,2
II	NEA	87,0	79,3	82,0	82,8
III	NCnt Cba	100,0	99,4	100,0	100,0
IV	S Cba	100,0	100,0	100,0	100,0
V	NCnt SFe	100,0	97,5	100,0	99,4
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	98,0	100,0	100,0	99,6
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	98,0	91,3	100,0	99,6
XI	SW BA-S LP	100,0	79,5	87,5	99,0
XII	SE BA	100,0	83,2	90,7	99,0
XIII	SL	100,0	100,0	100,0	100,0
XIV	Cuenca Sal	94,0	83,2	83,2	98,8
XV	Others	100,0	97,1	97,1	100,0
National		99,0	93,2	95,3	98,0

Soybean Condition: as of January 15, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	43,6	40,2	16,2
II	NEA	-	-	32,5	46,8	20,7
III	NCnt Cba	-	0,7	38,0	57,4	4,0
IV	S Cba	-	-	41,6	54,7	3,7
V	NCnt SFe	-	-	10,6	66,0	23,4
VI	North Belt	-	-	30,0	53,2	16,8
VII	South Belt	-	-	17,4	74,3	8,3
VIII	ECnt ER	-	6,8	53,0	31,3	9,0
IX	N LP-W BA	-	-	51,5	47,1	1,4
X	Cnt BA	-	4,2	77,0	18,8	-
XI	SW BA-S LP	-	5,1	40,7	54,1	-
XII	SE BA	-	-	34,5	60,8	4,7
XIII	SL	-	0,9	77,1	22,0	-
XIV	Cuenca Sal	-	3,7	63,6	32,7	-
XV	Others	-	-	23,9	69,5	6,6
National		-	0,9	37,2	52,7	9,2
Nat. Jan 16, 2019		2,0	7,9	38,8	40,6	10,6

CROP CONDITION & DEVELOPMENT STAGES

Soybean Moisture Condition: as of January 15, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	5,3	43,6	51,1	-
II	NEA	-	-	20,5	79,5	-
III	NCnt Cba	-	1,4	64,4	34,2	-
IV	S Cba	-	9,1	75,2	15,0	0,7
V	NCnt SFe	-	1,5	49,2	43,9	5,4
VI	North Belt	-	-	46,2	51,5	2,3
VII	South Belt	-	1,3	66,4	31,3	1,0
VIII	ECnt ER	-	-	47,3	51,3	1,5
IX	N LP-W BA	-	5,9	68,8	25,3	-
X	Cnt BA	-	20,8	67,6	11,6	-
XI	SW BA-S LP	1,2	24,7	55,7	18,4	-
XII	SE BA	-	10,0	59,2	30,8	-
XIII	SL	2,3	22,7	60,8	14,2	-
XIV	Cuenca Sal	-	13,7	58,0	28,3	-
XV	Others	-	-	34,8	62,3	2,9
National		0,1	4,5	56,8	37,7	1,0
Nat. Jan 16, 2019		-	0,8	35,7	46,4	17,2

Soybean Leaf Development: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	91,6	52,8	69,5	92,4
IV	S Cba	93,5	73,8	73,8	94,5
V	NCnt SFe	88,9	62,6	62,6	90,0
VI	North Belt	96,9	65,1	65,1	95,4
VII	South Belt	94,1	73,5	73,5	95,5
VIII	ECnt ER	94,7	75,0	75,0	96,5
IX	N LP-W BA	95,1	72,4	72,4	94,3
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		85,6	59,7	62,7	85,3

CROP CONDITION & DEVELOPMENT STAGES

Soybean Beginning Bloom: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	45,2	33,3	38,9	59,9
IV	S Cba	49,0	42,8	52,4	58,8
V	NCnt SFe	36,1	31,3	37,6	56,3
VI	North Belt	54,4	44,3	54,1	65,6
VII	South Belt	52,5	51,5	61,8	68,6
VIII	ECnt ER	38,7	35,3	42,0	61,9
IX	N LP-W BA	47,4	40,5	55,0	50,4
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		37,4	31,5	39,9	48,6

Soybean Beginning Pod: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	19,1	6,9	19,4	27,3
IV	S Cba	16,7	13,3	19,9	26,8
V	NCnt SFe	7,9	3,1	8,8	25,0
VI	North Belt	19,2	7,8	20,2	26,0
VII	South Belt	13,6	4,4	21,3	21,9
VIII	ECnt ER	3,9	-	-	15,0
IX	N LP-W BA	12,9	7,2	13,7	13,0
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		10,5	4,8	12,1	16,6

CROP CONDITION & DEVELOPMENT STAGES

Soybean Beginning Seed: as of January 15, 2020

Regions	January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I NWA	-	-	-	-
II NEA	-	-	-	-
III NCnt Cba	-	-	-	2,7
IV S Cba	-	-	-	-
V NCnt SFe	-	1,3	3,1	-
VI North Belt	-	-	-	1,4
VII South Belt	-	-	-	0,9
VIII ECnt ER	-	-	-	-
IX N LP-W BA	-	-	-	-
X Cnt BA	-	-	-	-
XI SW BA-S LP	-	-	-	-
XII SE BA	-	-	-	-
XIII SL	-	-	-	-
XIV Cuenca Sal	-	-	-	-
XV Others	-	-	-	-
National	-	0,1	0,2	0,6

Early Corn

Early Corn Planted: as of January 15, 2020

Regions	January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I NWA	65,0	30,0	45,0	56,4
II NEA	42,0	37,0	51,0	59,0
III NCnt Cba	100,0	100,0	100,0	100,0
IV S Cba	100,0	100,0	100,0	100,0
V NCnt SFe	100,0	100,0	100,0	100,0
VI North Belt	100,0	100,0	100,0	100,0
VII South Belt	100,0	100,0	100,0	100,0
VIII ECnt ER	100,0	100,0	100,0	100,0
IX N LP-W BA	100,0	100,0	100,0	100,0
X Cnt BA	100,0	100,0	100,0	100,0
XI SW BA-S LP	100,0	100,0	100,0	100,0
XII SE BA	100,0	100,0	100,0	100,0
XIII SL	100,0	100,0	100,0	100,0
XIV Cuenca Sal	100,0	100,0	100,0	100,0
XV Others	100,0	35,0	50,0	100,0
National	84,0	83,1	86,8	91,2

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Condition: as of January 15, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	60,0	40,0	-
II	NEA	-	-	60,0	35,0	5,0
III	NCnt Cba	-	3,0	50,0	43,0	4,0
IV	S Cba	-	5,0	55,0	40,0	-
V	NCnt SFe	-	-	50,0	45,0	5,0
VI	North Belt	-	3,0	29,0	60,0	8,0
VII	South Belt	-	3,0	35,0	54,0	8,0
VIII	ECnt ER	-	2,0	55,0	43,0	-
IX	N LP-W BA	-	3,0	35,0	60,0	2,0
X	Cnt BA	-	3,0	45,0	52,0	-
XI	SW BA-S LP	5,0	40,0	50,0	5,0	-
XII	SE BA	-	5,0	50,0	45,0	-
XIII	SL	-	10,0	60,0	30,0	-
XIV	Cuenca Sal	-	10,0	40,0	50,0	-
XV	Others	-	5,0	55,0	40,0	-
National		0,2	4,1	43,8	48,2	3,8
Nat. Jan 16, 2019		0,1	4,2	33,7	48,2	13,8

Early Corn Moisture Condition: as of January 15, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	10,0	50,0	40,0	-
II	NEA	-	5,0	50,0	45,0	-
III	NCnt Cba	-	3,0	47,0	50,0	-
IV	S Cba	-	8,0	50,0	42,0	-
V	NCnt SFe	-	15,0	40,0	40,0	5,0
VI	North Belt	-	9,0	46,0	40,0	5,0
VII	South Belt	-	8,0	39,0	50,0	3,0
VIII	ECnt ER	-	-	40,0	60,0	-
IX	N LP-W BA	-	1,0	54,0	45,0	-
X	Cnt BA	-	8,0	50,0	42,0	-
XI	SW BA-S LP	10,0	50,0	30,0	10,0	-
XII	SE BA	5,0	20,0	50,0	25,0	-
XIII	SL	-	-	60,0	40,0	-
XIV	Cuenca Sal	3,0	20,0	40,0	37,0	-
XV	Others	-	-	60,0	40,0	-
National		0,6	8,4	46,5	43,0	1,5
Nat. Jan 16, 2019		-	2,2	38,9	47,7	11,2

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Leaf Development: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	49,0	14,0	30,0	30,2
II	NEA	35,0	15,0	39,0	24,0
III	NCnt Cba	100,0	100,0	100,0	91,4
IV	S Cba	100,0	100,0	100,0	96,2
V	NCnt SFe	100,0	100,0	100,0	92,0
VI	North Belt	100,0	100,0	100,0	97,8
VII	South Belt	100,0	100,0	100,0	98,0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100,0	100,0	100,0	99,6
X	Cnt BA	100,0	100,0	100,0	100,0
XI	SW BA-S LP	-	-	-	-
XII	SE BA	100,0	100,0	100,0	99,4
XIII	SL	100,0	100,0	100,0	98,0
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		82,9	78,1	83,4	77,2

Early Corn Pollination: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	0,6
II	NEA	-	-	-	0,6
III	NCnt Cba	100,0	97,0	100,0	82,8
IV	S Cba	100,0	100,0	100,0	89,8
V	NCnt SFe	100,0	100,0	100,0	91,4
VI	North Belt	100,0	100,0	100,0	96,8
VII	South Belt	100,0	100,0	100,0	97,6
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100,0	91,0	100,0	89,0
X	Cnt BA	100,0	89,0	100,0	90,8
XI	SW BA-S LP	-	-	-	-
XII	SE BA	42,0	25,0	55,0	47,8
XIII	SL	91,0	65,0	90,0	85,0
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		68,4	67,8	71,8	62,8

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Silking: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	100,0	84,0	95,0	78,2
IV	S Cba	100,0	95,0	100,0	83,2
V	NCnt SFe	100,0	100,0	100,0	91,4
VI	North Belt	100,0	100,0	100,0	95,0
VII	South Belt	100,0	96,0	100,0	95,6
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	99,0	83,0	100,0	86,4
X	Cnt BA	95,0	57,0	89,0	82,8
XI	SW BA-S LP	-	-	-	-
XII	SE BA	32,0	16,0	40,0	32,0
XIII	SL	82,0	52,0	85,0	73,8
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		67,2	61,7	69,4	58,6

Early Corn Grain Growth: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	22,0	12,0	23,0	17,6
IV	S Cba	14,0	6,0	20,0	18,2
V	NCnt SFe	85,0	57,0	83,0	71,0
VI	North Belt	68,0	30,0	65,0	52,4
VII	South Belt	26,0	14,0	40,0	32,2
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	9,0	5,0	10,0	15,2
X	Cnt BA	6,0	1,0	8,0	9,4
XI	SW BA-S LP	-	-	-	0,4
XII	SE BA	-	-	-	-
XIII	SL	6,0	1,0	3,0	1,2
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		24,8	11,9	25,6	20,0

CROP CONDITION & DEVELOPMENT STAGES

Early Corn Mature: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	2,0	-	-	0,4
IV	S Cba	-	-	-	0,4
V	NCnt SFe	25,0	8,0	15,0	20,4
VI	North Belt	6,0	-	1,0	2,8
VII	South Belt	-	-	1,0	1,4
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		3,0	0,5	1,1	1,8

Late Corn

Late Corn Planted: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	98,0	100,0	100,0	98,6
IV	S Cba	100,0	100,0	100,0	99,6
V	NCnt SFe	50,0	56,0	70,0	70,4
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	86,0	80,0	90,0	92,0
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	100,0	100,0	100,0	100,0
XI	SW BA-S LP	100,0	100,0	100,0	100,0
XII	SE BA	100,0	100,0	100,0	100,0
XIII	SL	100,0	100,0	100,0	99,0
XIV	Cuenca Sal	100,0	100,0	100,0	100,0
XV	Others	-	-	-	-
National		96,0	96,7	97,9	91,4

CROP CONDITION & DEVELOPMENT STAGES

Late Corn Condition: as of January 15, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	-	-	-
II	NEA	-	-	-	-	-
III	NCnt Cba	-	-	38,0	52,0	10,0
IV	S Cba	-	-	35,0	50,0	15,0
V	NCnt SFe	-	-	20,0	68,0	12,0
VI	North Belt	-	-	25,0	65,0	10,0
VII	South Belt	-	-	65,0	27,0	8,0
VIII	ECnt ER	-	-	50,0	50,0	-
IX	N LP-W BA	-	-	40,0	50,0	10,0
X	Cnt BA	-	4,0	35,0	57,0	4,0
XI	SW BA-S LP	5,0	12,0	48,0	35,0	-
XII	SE BA	-	4,0	62,0	34,0	-
XIII	SL	-	10,0	50,0	40,0	-
XIV	Cuenca Sal	-	15,0	35,0	40,0	10,0
XV	Others	-	-	-	-	-
National		0,1	2,0	39,6	49,4	8,9
Nat. Jan 16, 2019		0,6	4,8	46,5	41,0	7,1

Late Corn Moisture Condition: as of January 15, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	-	-	-	-
II	NEA	-	-	-	-	-
III	NCnt Cba	-	-	43,0	57,0	-
IV	S Cba	-	-	40,0	60,0	-
V	NCnt SFe	-	5,0	50,0	37,0	8,0
VI	North Belt	-	9,0	45,0	46,0	-
VII	South Belt	-	10,0	45,0	44,0	1,0
VIII	ECnt ER	-	5,0	44,0	51,0	-
IX	N LP-W BA	-	4,0	45,0	51,0	-
X	Cnt BA	-	8,0	49,0	43,0	-
XI	SW BA-S LP	12,0	45,0	43,0	-	-
XII	SE BA	-	30,0	37,0	33,0	-
XIII	SL	-	30,0	40,0	30,0	-
XIV	Cuenca Sal	-	15,0	55,0	30,0	-
XV	Others	-	-	-	-	-
National		0,3	7,2	42,8	49,4	0,3
Nat. Jan 16, 2019		0,2	5,5	47,8	37,7	8,8

CROP CONDITION & DEVELOPMENT STAGES

Late Corn Leaf Development: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	77,0	81,0	85,0	48,0
IV	S Cba	80,0	79,0	82,0	66,0
V	NCnt SFe	31,0	15,0	35,0	31,2
VI	North Belt	75,0	70,0	80,0	61,4
VII	South Belt	76,0	71,0	82,0	69,8
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	85,0	81,0	90,0	77,6
X	Cnt BA	82,0	75,0	87,0	74,2
XI	SW BA-S LP	-	-	-	-
XII	SE BA	73,0	70,0	84,0	63,8
XIII	SL	86,0	74,0	82,0	72,0
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		75,3	71,8	79,4	60,4

Late Corn Pollination: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	-	1,0	2,0	3,0
IV	S Cba	4,0	2,0	5,0	11,0
V	NCnt SFe	-	-	-	1,0
VI	North Belt	9,0	3,0	6,0	12,6
VII	South Belt	-	3,0	5,0	7,8
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	1,0	2,0	4,0	10,4
X	Cnt BA	-	-	2,0	8,6
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	2,0	5,0
XIII	SL	-	-	4,0	13,8
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		1,4	1,1	3,2	7,6

CROP CONDITION & DEVELOPMENT STAGES

Late Corn Silking: as of January 15, 2020

Regions	January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I NWA	-	-	-	-
II NEA	-	-	-	-
III NCnt Cba	-	-	1,0	1,0
IV S Cba	-	-	1,0	3,0
V NCnt SFe	-	-	-	-
VI North Belt	-	-	1,0	4,0
VII South Belt	-	-	2,0	3,6
VIII ECnt ER	-	-	-	-
IX N LP-W BA	-	-	1,0	1,2
X Cnt BA	-	-	-	-
XI SW BA-S LP	-	-	-	-
XII SE BA	-	-	-	1,0
XIII SL	-	-	1,0	5,0
XIV Cuenca Sal	-	-	-	-
XV Others	-	-	-	-
National	-	-	0,8	2,0

Corn

Corn Planted: as of January 15, 2020

Regions	January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I NWA	55,0	30,0	45,0	50,4
II NEA	38,0	37,0	51,0	56,2
III NCnt Cba	96,0	100,0	100,0	98,4
IV S Cba	98,0	100,0	100,0	99,4
V NCnt SFe	61,0	75,8	83,5	80,0
VI North Belt	99,0	100,0	100,0	99,8
VII South Belt	99,0	100,0	100,0	99,8
VIII ECnt ER	84,0	93,0	96,5	94,8
IX N LP-W BA	98,0	100,0	100,0	99,6
X Cnt BA	99,0	100,0	100,0	99,8
XI SW BA-S LP	98,0	100,0	100,0	99,6
XII SE BA	96,0	100,0	100,0	99,2
XIII SL	98,0	100,0	100,0	99,0
XIV Cuenca Sal	97,0	100,0	100,0	99,4
XV Others	100,0	35,0	50,0	92,6
National	86,0	88,2	91,0	89,8

CROP CONDITION & DEVELOPMENT STAGES

Corn Condition: as of January 15, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	60,0	40,0	-
II	NEA	-	-	60,0	35,0	5,0
III	NCnt Cba	-	0,7	40,9	49,8	8,6
IV	S Cba	-	1,3	40,2	47,4	11,1
V	NCnt SFe	-	-	36,2	55,6	8,2
VI	North Belt	-	2,8	28,8	60,3	8,1
VII	South Belt	-	2,8	37,1	52,1	8,0
VIII	ECnt ER	-	1,3	53,4	45,3	-
IX	N LP-W BA	-	2,0	36,8	56,5	4,8
X	Cnt BA	-	3,4	41,4	53,8	1,4
XI	SW BA-S LP	5,0	29,6	49,3	16,1	-
XII	SE BA	-	4,7	54,2	41,2	-
XIII	SL	-	10,0	52,6	37,4	-
XIV	Cuenca Sal	-	11,8	38,3	46,5	3,5
XV	Others	-	5,0	55,0	40,0	-
National		0,1	3,2	42,1	48,7	5,9
Nat. Jan 16, 2019		0,3	4,5	39,9	44,7	10,6

Corn Moisture Condition: as of January 15, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	10,0	50,0	40,0	-
II	NEA	-	5,0	50,0	45,0	-
III	NCnt Cba	-	0,7	44,0	55,3	-
IV	S Cba	-	2,1	42,6	55,3	-
V	NCnt SFe	-	10,4	44,6	38,6	6,4
VI	North Belt	-	9,0	45,9	40,4	4,7
VII	South Belt	-	8,1	39,4	49,6	2,9
VIII	ECnt ER	-	1,6	41,3	57,1	-
IX	N LP-W BA	-	2,1	50,9	47,1	-
X	Cnt BA	-	8,0	49,6	42,4	-
XI	SW BA-S LP	10,7	48,2	34,8	6,3	-
XII	SE BA	3,3	23,5	45,5	27,8	-
XIII	SL	-	22,2	45,2	32,6	-
XIV	Cuenca Sal	2,0	18,3	45,3	34,6	-
XV	Others	-	-	60,0	40,0	-
National		0,5	7,9	45,0	45,6	1,0
Nat. Jan 16, 2019		0,1	3,8	43,2	42,9	10,0

CROP CONDITION & DEVELOPMENT STAGES

Corn Leaf Development: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	49,0	14,0	30,0	29,2
II	NEA	35,0	15,0	39,0	27,0
III	NCnt Cba	80,0	85,6	88,6	56,6
IV	S Cba	84,0	84,5	86,7	76,2
V	NCnt SFe	55,0	53,3	64,3	57,4
VI	North Belt	97,0	98,2	98,8	93,2
VII	South Belt	96,0	98,0	98,7	94,0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	92,0	93,4	96,5	89,4
X	Cnt BA	90,0	91,0	95,3	85,8
XI	SW BA-S LP	-	-	-	-
XII	SE BA	86,0	89,5	94,4	90,0
XIII	SL	89,0	80,8	86,7	78,4
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		79,3	75,7	81,9	70,6

Corn Pollination: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	0,6
II	NEA	-	-	-	0,6
III	NCnt Cba	15,0	24,0	25,5	18,6
IV	S Cba	23,0	27,5	29,7	36,0
V	NCnt SFe	35,0	45,0	45,0	36,2
VI	North Belt	88,0	94,2	94,4	85,0
VII	South Belt	85,0	93,2	93,4	81,8
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	50,0	59,9	66,4	56,8
X	Cnt BA	45,0	57,0	64,7	52,2
XI	SW BA-S LP	-	-	-	-
XII	SE BA	21,0	16,3	36,5	37,0
XIII	SL	18,0	16,9	26,4	32,4
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		37,4	42,8	46,1	41,2

CROP CONDITION & DEVELOPMENT STAGES

Corn Silking: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	15,0	20,2	23,6	16,2
IV	S Cba	20,0	24,7	26,7	27,8
V	NCnt SFe	35,0	45,0	45,0	35,4
VI	North Belt	87,0	94,0	94,1	82,6
VII	South Belt	85,0	89,3	93,1	80,2
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	50,0	54,0	65,4	51,4
X	Cnt BA	43,0	36,5	57,0	43,2
XI	SW BA-S LP	-	-	-	-
XII	SE BA	16,0	10,4	26,0	23,8
XIII	SL	16,0	13,5	22,8	21,8
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		36,3	38,6	43,7	36,0

Corn Grain Growth: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	3,0	2,9	5,5	3,6
IV	S Cba	3,0	1,6	5,2	4,2
V	NCnt SFe	30,0	25,7	37,4	28,2
VI	North Belt	59,0	28,2	61,1	43,8
VII	South Belt	22,0	13,0	37,2	22,8
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	4,0	3,3	6,5	7,8
X	Cnt BA	3,0	0,6	5,1	3,8
XI	SW BA-S LP	-	-	-	2,0
XII	SE BA	-	-	-	-
XIII	SL	1,0	0,3	0,8	0,2
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		13,3	7,4	16,0	11,2

CROP CONDITION & DEVELOPMENT STAGES

Corn Mature: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	-	-	-	-
III	NCnt Cba	-	-	-	0,2
IV	S Cba	-	-	-	-
V	NCnt SFe	9,0	3,6	6,8	8,6
VI	North Belt	5,0	-	0,9	2,2
VII	South Belt	-	-	0,9	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		1,5	0,3	0,7	1,2

Sunflower

Sunflower Condition: as of January 15, 2020

Regions		Very Poor %	Poor %	Fair %	Good %	Excellent %
I	NWA	-	-	-	-	-
II	NEA	-	11,0	35,0	53,0	1,0
III	NCnt Cba	-	-	-	-	-
IV	S Cba	-	-	-	-	-
V	NCnt SFe	-	-	28,0	65,0	7,0
VI	North Belt	-	-	-	-	-
VII	South Belt	-	-	-	-	-
VIII	ECnt ER	-	-	-	-	-
IX	N LP-W BA	-	-	38,0	54,0	8,0
X	Cnt BA	-	-	-	-	-
XI	SW BA-S LP	-	21,0	60,0	19,0	-
XII	SE BA	-	2,0	47,0	51,0	-
XIII	SL	-	-	-	-	-
XIV	Cuenca Sal	-	2,0	35,0	52,0	11,0
XV	Others	-	-	-	-	-
National		-	7,8	44,3	45,1	2,8
Nat. Jan 16, 2019		2,6	13,1	31,3	41,8	11,3

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Moisture Condition: as of January 15, 2020

Regions		Dry %	Poor %	Favourable %	Optimum %	Wet %
I	NWA	-	-	-	-	-
II	NEA	-	-	29,0	61,0	10,0
III	NCnt Cba	-	-	-	-	-
IV	S Cba	-	-	-	-	-
V	NCnt SFe	-	-	41,0	57,0	2,0
VI	North Belt	-	-	-	-	-
VII	South Belt	-	-	-	-	-
VIII	ECnt ER	-	-	-	-	-
IX	N LP-W BA	-	-	39,0	61,0	-
X	Cnt BA	-	-	-	-	-
XI	SW BA-S LP	10,0	39,0	51,0	-	-
XII	SE BA	-	8,0	63,0	29,0	-
XIII	SL	-	-	-	-	-
XIV	Cuenca Sal	-	3,0	50,0	47,0	-
XV	Others	-	-	-	-	-
National		2,5	12,4	49,7	34,1	1,4
Nat. Jan 16, 2019		0,2	3,8	41,8	39,6	14,6

Sunflower Flowerbud Stage: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100,0	100,0	100,0	100,0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	100,0	100,0	100,0	100,0
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100,0	92,0	100,0	100,0
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	65,0	63,0	77,0	76,6
XII	SE BA	68,0	58,0	73,0	82,8
XIII	SL	-	-	-	-
XIV	Cuenca Sal	83,0	88,0	100,0	92,4
XV	Others	-	-	-	-
National		84,7	80,1	88,5	89,9

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Flowering Stage: as of January 15, 2020

Regions	January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I NWA	-	-	-	-
II NEA	100,0	100,0	100,0	100,0
III NCnt Cba	-	-	-	-
IV S Cba	-	-	-	-
V NCnt SFe	100,0	100,0	100,0	100,0
VI North Belt	-	-	-	-
VII South Belt	-	-	-	-
VIII ECnt ER	-	-	-	-
IX N LP-W BA	61,0	61,0	72,0	68,4
X Cnt BA	-	-	-	-
XI SW BA-S LP	33,0	30,0	44,0	43,0
XII SE BA	47,0	32,0	49,0	54,4
XIII SL	-	-	-	-
XIV Cuenca Sal	77,0	45,0	61,0	58,4
XV Others	-	-	-	-
National	69,4	60,5	70,0	69,1

Sunflower Seed Development: as of January 15, 2020

Regions	January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I NWA	-	-	-	-
II NEA	100,0	100,0	100,0	100,0
III NCnt Cba	-	-	-	-
IV S Cba	-	-	-	-
V NCnt SFe	100,0	100,0	100,0	100,0
VI North Belt	-	-	-	-
VII South Belt	-	-	-	-
VIII ECnt ER	-	-	-	-
IX N LP-W BA	32,0	7,0	21,0	27,0
X Cnt BA	-	-	-	-
XI SW BA-S LP	-	-	-	6,4
XII SE BA	3,0	-	-	8,4
XIII SL	-	-	-	-
XIV Cuenca Sal	28,0	-	16,0	24,6
XV Others	-	-	-	-
National	45,8	36,2	39,1	42,0

CROP CONDITION & DEVELOPMENT STAGES

Sunflower Mature: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	100,0	100,0	100,0	100,0
III	NCnt Cba	-	-	-	-
IV	S Cba	-	-	-	-
V	NCnt SFe	89,0	85,0	100,0	89,8
VI	North Belt	-	-	-	-
VII	South Belt	-	-	-	-
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	-	-	-	0,4
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		37,2	31,6	34,1	31,9

Sunflower Harvested: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	-	-	-	-
II	NEA	68,0	43,3	53,7	76,0
III	NCnt Cba	23,0	-	29,0	34,6
IV	S Cba	2,0	-	-	1,4
V	NCnt SFe	34,0	12,7	36,7	32,4
VI	North Belt	12,0	-	-	11,4
VII	South Belt	-	-	-	1,2
VIII	ECnt ER	-	-	-	0,4
IX	N LP-W BA	-	-	-	-
X	Cnt BA	-	-	-	-
XI	SW BA-S LP	-	-	-	-
XII	SE BA	-	-	-	-
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	8,0	-	-	3,6
National		20,5	9,9	15,4	17,5

CROP CONDITION & DEVELOPMENT STAGES

Wheat


Wheat Harvested: as of January 15, 2020

Regions		January 16, 2019 %	January 8, 2020 %	January 15, 2020 %	2014/2019 Average %
I	NWA	100,0	100,0	100,0	100,0
II	NEA	100,0	100,0	100,0	100,0
III	NCnt Cba	100,0	100,0	100,0	100,0
IV	S Cba	100,0	100,0	100,0	100,0
V	NCnt SFe	100,0	100,0	100,0	100,0
VI	North Belt	100,0	100,0	100,0	100,0
VII	South Belt	100,0	100,0	100,0	100,0
VIII	ECnt ER	-	-	-	-
IX	N LP-W BA	100,0	100,0	100,0	100,0
X	Cnt BA	100,0	100,0	100,0	100,0
XI	SW BA-S LP	100,0	100,0	100,0	97,0
XII	SE BA	100,0	54,2	100,0	94,0
XIII	SL	-	-	-	-
XIV	Cuenca Sal	-	-	-	-
XV	Others	-	-	-	-
National		100,0	95,5	100,0	98,9

Wheat 2019/20 Development in Argentina

CROP CONDITION & DEVELOPMENT STAGES

Crop Condition: ■ Poor/Very poor ■ Fair ■ Excellent/Good
 Development Stages: — Planting — Leaves Differentiation — Tillering — Steam Extension
— Flowering — Ripening — Mature - - - Harvest



Moisture Condition: ■ Dry/Poor ■ Favourable/Optimum ■ Wet

Source: Department of Agricultural Estimates - Buenos Aires Grain Exchange

As of Jan 16, 2020